

Weak stuffle algebras

Cécile Mammez

Univ. Lille, UMR 8524 - Laboratoire Paul Painlevé, F-59000 Lille, France

CNRS, UMR 8524, F-59000 Lille, France

e-mail : cecile.mammez@univ-lille.fr

Abstract. Motivated by q -shuffle products determined by Singer from q -analogues of multiple zeta values, we build in this article a generalisation of the shuffle and stuffle products in terms of weak shuffle and stuffle products. Then, we characterise weak shuffle products and give as examples the case of an alphabet of cardinality two or three. We focus on a comparison between algebraic structures respected in the classical case and in the weak case. As in the classical case, each weak shuffle product can be equipped with a dendriform structure. However, they have another behaviour towards the quadri-algebra and the Hopf algebra structure. We give some relations satisfied by weak stuffle products.

Keywords. Shuffle algebras, stuffle algebras, dendriform algebras, quadri-algebras, Hopf algebras;

Résumé. A partir de q -analogues aux fonctions zêta multiples, Singer détermine des q -battages. Ceci motive, dans cet article, la construction d'une généralisation des produits de battage et de battage contractant en produits de battage faibles et produits de battage contractant faibles. Nous caractérisons ensuite les battages faibles et donnons comme exemple le cas d'un alphabet à deux ou trois lettres. Nous comparons les structures algébriques respectées dans le cas classique et dans le cas faible. Comme dans le cas classique, tout battage faible peut être muni d'une structure d'algèbre dendriforme. En revanche, ils se comportent différemment face à la structure de quadri-algèbre et d'algèbre de Hopf. Nous donnons des relations vérifiées par les battages contractants faibles.

Mots-clés. Algèbres de battage, algèbres de battage contractant, algèbres dendriformes, quadri-algèbres, algèbres de Hopf.

AMS classification. 05A05, 05E40, 16T30, 68R15

Introduction

The notion of shuffle and stuffle algebras is widely used in several fields of mathematics. Indeed, they participate in the study of Rota-Baxter algebras with the notion of mixable shuffle algebras [14, 6, 20], in the study of Yang-Baxter algebras [21], in the study of quasi-symmetric functions and words algebras [13, 24, 25, 5, 4, 12, 33, 26], in the study of multiple zeta values [34, 15, 16, 19, 18, 17, 30, 8, 7] ...

The classical stuffle product comes from the product of classical multiple zeta values and is defined by the relation

$$au \square bv = a(u \square bv) + b(au \square v) + (a \diamond b)(u \square v)$$

where a and b are letters, u and v are words and \diamond is an associative and commutative product which is equal to 0 in the case of the classical shuffle product. Thus, the shuffle part of the

relation is symmetric and does not depend on letters of any words in the product. In his work, Singer focuses on q -shuffle products coming from q -analogs of multiples zeta values. This case enables the existence of some letters p and y satisfying a relation in the form of

$$yu \square pv = pv \square yu = y(u \square pv)$$

for any words u and v . This new q -shuffle relation is not symmetric and depends on the beginning of each word in the product. This leads to focus on new generalisations of shuffle and stuffle products [31, 7, 8].

In this article, we present a new generalisation of shuffle and stuffle algebras, we study their algebraic structures and compare them to the classical case. The article is organised as follows.

- In Section 1, we recall the classical notion of shuffle and stuffle product thanks to the multiple zeta values as well as the calculation by Singer of q -shuffle associated to the Schlesinger-Zudilin model and the Bradley-Zhao model.
- In Section 2, we define a generalisation of the classical shuffle product and the classical stuffle product called weak shuffle products and weak stuffle products and prove a characterisation of weak shuffle products. We detail the case of an alphabet of cardinality 2 or 3.
- In Section 3, we focus on algebraic structures respected by the classical shuffle product and we determine if the weak shuffle products respect them too. Thus we prove that weak shuffle products are dendriform but there are obstacles to the quadri-algebra structure.
- In Section 4, we express some relations satisfied by weak stuffle products and we express the q -shuffle given by Singer in terms of weak stuffle product. Besides, in the case of an infinite, countable and totally ordered alphabet $\{x_1, \dots, x_n, \dots\}$, we prove that, if the contracting part in the weak stuffle products is expressed as $f_3(x_i \otimes x_j) \in \mathbb{K}^* x_{i+j}$, then the shuffle part is the null product or the classical shuffle product. We give some informations more about weak stuffle products in the case of an alphabet of cardinality 2 or 3.
- In Section 5, we prove that a weak stuffle product is compatible with the deconcatenation coproduct if and only if the underlying weak shuffle product is the classical shuffle product and the contracting part is associative and commutative.
- Computation programs used to prove Lemma 18 are detailed in Section 6.

Acknowledgment: I would like to thank the anonymous referees for their useful comments and suggestions. I would like to thank all people who supported me for this work. This work was funding by the Laboratoire Paul Painlevé at Université de Lille, the Fédération de Recherche Mathématique des Hauts-de-France, the ANR Alcohol project ANR-19-CE40-0006 and the Labex CEMPI ANR-11-LABX-0007-01.

1 Reminders

1.1 Classical shuffle and stuffle algebras

We recall here the definition of the stuffle product in the context of the multiple zeta values.

Definition 1. Let s be an integer and let (k_1, \dots, k_s) be an s -tuple in $\mathbb{N}_{\geq 2} \times \mathbb{N}^{s-1}$. The multiple zeta value associated to (k_1, \dots, k_s) is

$$\zeta(k_1, \dots, k_s) = \sum_{\substack{(m_1, \dots, m_s) \in \mathbb{N} \\ m_1 > \dots > m_s > 0}} \frac{1}{m_1^{k_1} \dots m_s^{k_s}}.$$

On multiple zeta values, we consider the product of functions taking values in \mathbb{C} . For instance,

$$\begin{aligned}\zeta(n)\zeta(m) &= \zeta(m, n) + \zeta(n, m) + \zeta(m+n), \\ \zeta(n, p)\zeta(m) &= \zeta(m, n, p) + \zeta(n, m, p) + \zeta(n, p, m) + \zeta(n+m, p) + \zeta(n, p+m).\end{aligned}$$

Then, it leads to the following algebraic definition and following theorem [15].

Theorem 2. *Let $X = \{x_1, \dots, x_n, \dots\}$ be a countable alphabet. Let $\mathbb{K}\langle X \rangle$ be the algebra of words on the alphabet X . We define the product \star , called the stuffle product, by:*

$$\begin{aligned}u \star 1 &= 1 \star u = 1, \\ u \star 0 &= 0 \star u = 0, \\ x_i u \star x_j v &= x_i(u \star x_j v) + x_j(x_i u \star v) + x_{i+j}(u \star v)\end{aligned}$$

for any letters x_i and x_j and any words u and v .

Then

$$\begin{aligned}x_i u x_k \star x_j v x_l &= x_i(u x_k \star x_j v x_l) + x_j(x_i u x_k \star v x_l) + x_{i+j}(u x_k \star v x_l) \\ &= (x_i u \star x_j v x_l) x_k + (x_i u x_k \star x_j v) x_l + (x_i u \star x_j v) x_{k+l}\end{aligned}$$

and $(\mathbb{K}\langle X \rangle, \star)$ is an associative and commutative algebra.

It is possible to define another algebra:

Theorem 3. *Let $X = \{x_1, \dots, x_n, \dots\}$ be a countable alphabet. Let $\mathbb{K}\langle X \rangle$ be the algebra of words on the alphabet X . We define the product \sqcup , called the shuffle product, by:*

$$\begin{aligned}u \sqcup 1 &= 1 \sqcup u = 1, \\ u \sqcup 0 &= 0 \sqcup u = 0, \\ x_i u \sqcup x_j v &= x_i(u \sqcup x_j v) + x_j(x_i u \sqcup v)\end{aligned}$$

for any letters x_i and x_j and any words u and v .

Then

$$\begin{aligned}x_i u x_k \sqcup x_j v x_l &= x_i(u x_k \sqcup x_j v x_l) + x_j(x_i u x_k \sqcup v x_l) \\ &= (x_i u \sqcup x_j v x_l) x_k + (x_i u x_k \sqcup x_j v) x_l\end{aligned}$$

and $(\mathbb{K}\langle X \rangle, \sqcup)$ is an associative and commutative algebra.

Theorem 4. *Let $X = \{x_1, \dots, x_n, \dots\}$ be a countable alphabet. The algebras $(\mathbb{K}\langle X \rangle, \star)$ and $(\mathbb{K}\langle X \rangle, \sqcup)$ are isomorphic.*

Proof. This theorem was proved by Hoffman [16, Theorem 2.5] by describing an explicit isomorphism \exp . Another construction of \exp leading to the proof of this theorem is given in [26, Proposition 41]. \square

1.2 q -shuffle products for the Schlesinger-Zudilin model and the Bradley-Zhao model.

Let q be real number such that $0 < q < 1$. A q -analogue of a positive integer m is defined by

$$[m]_q = \frac{1 - q^m}{1 - q} = 1 + q + \dots + q^{m-1}.$$

The Schlesinger-Zudilin model [28, 36] is defined as the following q -sum:

$$\zeta_q^{SZ}(k_1, \dots, k_n) = (1 - q)^{-(k_1 + \dots + k_n)} \sum_{\substack{(m_1, \dots, m_s) \in \mathbb{N} \\ m_1 > \dots > m_s > 0}} \frac{q^{m_1 k_1 + \dots + m_n k_n}}{[m_1]_q^{k_1} \dots [m_n]_q^{k_n}}$$

$$= \sum_{\substack{(m_1, \dots, m_s) \in \mathbb{N} \\ m_1 > \dots > m_s > 0}} \frac{q^{m_1 k_1 + \dots + m_n k_n}}{(1 - q^{m_1})^{k_1} \dots (1 - q^{m_n})^{k_n}}$$

for any $(k_1, \dots, k_n) \in (\mathbb{N}^*)^n$.

The Bradley-Zhao model [2, 35] is defined as the following q -sum:

$$\begin{aligned} \zeta_q^{BZ}(k_1, \dots, k_n) &= (1 - q)^{-(k_1 + \dots + k_n)} \sum_{\substack{(m_1, \dots, m_s) \in \mathbb{N} \\ m_1 > \dots > m_s > 0}} \frac{q^{m_1(k_1-1) + \dots + m_n(k_n-1)}}{[m_1]_q^{k_1} \dots [m_n]_q^{k_n}} \\ &= \sum_{\substack{(m_1, \dots, m_s) \in \mathbb{N} \\ m_1 > \dots > m_s > 0}} \frac{q^{m_1(k_1-1) + \dots + m_n(k_n-1)}}{(1 - q^{m_1})^{k_1} \dots (1 - q^{m_n})^{k_n}} \end{aligned}$$

for any $(k_1, \dots, k_n) \in \mathbb{N}^n$ with $k_1 \geq 2$.

From those two models, Singer defined two q -shuffle products corresponding to the algebraic version of the Schlesinger-Zudilin model and the Bradley-Zhao model and proved the following two theorems in [29, 30, 31]:

Theorem 5 (Singer). *Let $X = \{y, p\}$ be an alphabet. The q -shuffle product associated to the Schlesinger-Zudilin model is given by: for any words u and v ,*

1. $1 \sqcup_{SZ} u = u \sqcup_{SZ} 1 = u$,
2. $yu \sqcup_{SZ} v = v \sqcup_{SZ} yu = y(u \sqcup_{SZ} v)$,
3. $pu \sqcup_{SZ} pv = p(u \sqcup_{SZ} pv) + p(pu \sqcup_{SZ} v) + p(u \sqcup_{SZ} v)$.

Besides, it is an associative and commutative product.

Theorem 6 (Singer). *Let $X = \{y, p, \bar{p}\}$ be an alphabet. The q -shuffle product associated to the Bradley-Zhao model is given by: for any words u and v ,*

1. $1 \sqcup_{BZ} u = u \sqcup_{BZ} 1 = u$,
2. $yu \sqcup_{BZ} v = v \sqcup_{BZ} yu = y(u \sqcup_{BZ} v)$,
3. $au \sqcup_{BZ} bv = a(u \sqcup_{BZ} bv) + b(au \sqcup_{BZ} v) + [a, b]a(u \sqcup_{BZ} v)$ where

$$a, b \in \{p, \bar{p}\}, [p, p] = -[\bar{p}, \bar{p}] = 1 \text{ and } [p, \bar{p}] = [\bar{p}, p] = 0.$$

Besides, it is an associative and commutative product.

2 Definition and characterisation of weak shuffle products

The aim of this section is to define a generalisation of the classical shuffle product, the classical stuffle product, and the two q -shuffle products given by the Schlesinger-Zudilin model and the Bradley-Zhao model. We give and prove a characterisation of weak shuffle products too. Then we explicit the case of an alphabet of cardinality 2 or 3.

2.1 Characterisation

Definition 7. *An alphabet is a non-empty finite or countable set X .*

Definition 8. *Let X be an alphabet. We denote by X^* the set of words on the alphabet X and by $\mathbb{K}\langle X \rangle$ the tensor algebra generated by X (i.e. the algebra of words on X). The space $\mathbb{K}\langle X \rangle$ is graded by the length of words.*

Definition 9. Let X be an alphabet. A weak stuffle product on $\mathbb{K}\langle X \rangle$ is an associative and commutative product \square such that for any $(a, b) \in (X)^2$ and any $(u, v) \in (X^*)^2$

$$\begin{aligned} u\square 1 &= 1\square u = u, \\ u\square 0 &= 0\square u = 0, \\ au\square bv &= f_1(a \otimes b)a(u\square bv) + f_2(a \otimes b)b(au\square v) + f_3(a \otimes b)(u\square v) \end{aligned}$$

where

1. f_1 and f_2 are linear maps from $\mathbb{K}.X \otimes \mathbb{K}.X$ to \mathbb{K} ,
2. $f_3 = kg$ is a linear map from $\mathbb{K}.X \otimes \mathbb{K}.X$ to $\mathbb{K}.X$ such that $k(a \otimes b) \in \mathbb{K}$ and $g(a \otimes b) \in X$ for any $(a, b) \in X^2$,
3. If $f_3 \equiv 0$ then the product \square is called a weak shuffle product.

Examples. Let $X = \{x_1, \dots, x_n, \dots\}$ be an infinite alphabet.

1. The classical shuffle product on $\mathbb{K}\langle X \rangle$ is a weak stuffle product where $f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any $(a, b) \in X^2$, and $f_3 \equiv 0$.
2. The classical stuffle product on $\mathbb{K}\langle X \rangle$ is a weak stuffle product where $f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any $(a, b) \in X^2$, and $f_3(x_i \otimes x_j) = x_{i+j}$ for any $(i, j) \in (\mathbb{N}^*)^2$.
3. The stuffle products on $\mathbb{K}\langle X \rangle$ given by Hoffman and Ihara in [18] is a weak stuffle product where $f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any $(a, b) \in X^2$, and $f_3(x_i \otimes x_j) = -x_{i+j}$ for any $(i, j) \in (\mathbb{N}^*)^2$.

Theorem 10. Let \square be a product on $\mathbb{K}\langle X \rangle$. The map \square is a weak shuffle product if and only if, for any distinct letters a, b , and c in X :

1. $f_1(a \otimes b) = f_2(b \otimes a)$.
2. (a) either $f_1(a \otimes a) = f_2(a \otimes a) = \alpha$ with $\alpha \in \{0, 1\}$ and
 - i. $f_1(a \otimes b)f_1(b \otimes a)[f_1(a \otimes a) - 1] = 0$,
 - ii. $f_1(a \otimes a)f_1(a \otimes b)[f_1(a \otimes b) - 1] = 0$,
 - iii. $f_1(a \otimes a)f_1(b \otimes a)[f_1(b \otimes a) - 1] = 0$.
 (b) or $f_1(a \otimes a) = \alpha$, $f_2(a \otimes a) = 1 - \alpha$ with $\alpha \in \mathbb{R}$ and
 - i. $f_1(a \otimes b) = 1$,
 - ii. $f_1(b \otimes a) = 0$.
3. $f_1(a \otimes b)f_1(b \otimes c)[f_1(a \otimes c) - 1] = 0$.
4. $f_3 \equiv 0$.

Remark. It is sometimes useful to use in calculations the following statement induced by the item 2b of the Theorem 10:

"If $f_1(a \otimes b) = 0$ or $f_1(b \otimes a) \neq 0$ then $f_1(a \otimes a) = f_2(a \otimes a) = \alpha$ with $\alpha \in \{0, 1\}$."

Proof. Let us prove first the direct implication. Let us assume \square is a weak shuffle product. Let a, b , and c be three distinct letters. Then, by direct calculations,

(A) $a\square b = b\square a$ gives relation $f_1(a \otimes b) = f_2(b \otimes a)$.

(B) $a\square aa = aa\square a$ gives $f_1(a \otimes a) = f_2(a \otimes a)$ or $f_1(a \otimes a) = 1 - f_2(a \otimes a)$.

- (C) $a \square ab = ab \square a$ gives, if $f_1(a \otimes b) = 0$ or $f_1(b \otimes a) \neq 0$, that $f_1(a \otimes a) = f_2(a \otimes a)$. Thus, if $f_1(a \otimes a) = 1 - f_2(a \otimes a)$ and $f_1(a \otimes a) \neq \frac{1}{2}$ then $f_1(a \otimes b) \neq 0$ and $f_1(b \otimes a) = 0$. The relation $a \square ab = ab \square a$ implies $f_1(a \otimes b) = 1$.
- (D) $(a \square a) \square b = a \square (a \square b) = (a \square b) \square a$ with $f_1(a \otimes a) = f_2(a \otimes a)$ give
- (a) $f_1(a \otimes b)f_1(b \otimes a)[f_1(a \otimes a) - 1] = 0$,
 - (b) $f_1(a \otimes a)f_1(a \otimes b)[f_1(a \otimes b) - 1] = 0$,
 - (c) $f_1(a \otimes a)f_1(b \otimes a)[f_1(b \otimes a) - 1] = 0$.
- (E) $(a \square b) \square c = a \square (b \square c)$ gives $f_1(a \otimes b)f_1(b \otimes c)[f_1(a \otimes c) - 1] = 0$.
- (F) $(a \square a) \square ab = a \square (a \square ab)$ implies that if $f_1(a \otimes a) = 1 - f_2(a \otimes a) = \frac{1}{2}$ then $f_1(a \otimes b) = 1$ and $f_1(b \otimes a) = 0$.
- (G) $(a \square a) \square aa = a \square (a \square aa)$ and $(a \square a) \square aaa = a \square (a \square aaa)$ implies that if $f_1(a \otimes a) = f_2(a \otimes a) = \alpha$ then $\alpha \in \{0, 1, \frac{1}{2}\}$.
- (H) Cases $ba \square a = a \square ba$, $aa \square b = b \square aa$, $ab \square c = c \square ab$ and $(a \square a) \square a = a \square (a \square a)$ do not give any further relations.

As consequences, in the theorem 10,

- the item 1 is proved by the item (A),
- the item 2a is proved by the items (B), (D), (F) and (G),
- the item 2b is proved by the items (B), (C) and (F),
- the item 3 is proved by the item (E),
- the item 4 is satisfied by the definition of a weak shuffle product.

Conversly, if \square satisfies all relations given in Theorem 10 then for any couple (u, v) and any triple (w_1, w_2, w_3) of words such that $\text{length}(u) + \text{length}(v) \leq 3$ and $\text{length}(w_1) + \text{length}(w_2) + \text{length}(w_3) \leq 3$ one has: $u \square v = v \square u$ and $(w_1 \square w_2) \square w_3 = w_1 \square (w_2 \square w_3)$.

We assume now there exists an integer $n \geq 3$ such that $u \square v = v \square u$ and $(w_1 \square w_2) \square w_3 = w_1 \square (w_2 \square w_3)$ for any words u, v, w_1, w_2 with $\text{length}(u) + \text{length}(v) \leq n$ and $\text{length}(w_1) + \text{length}(w_2) + \text{length}(w_3) \leq n$.

Let now u and v be two words such that $\text{length}(u) + \text{length}(v) = n + 1$. Then there exist two letters a and b and two words w_1 and w_2 (not necessarily non-empty) such that $u = aw_1$ and $v = bw_2$. Then, by induction, we get:

case $a \neq b$.

$$\begin{aligned} u \square v &= f_1(a \otimes b)a(w_1 \square bw_2) + f_1(b \otimes a)b(aw_1 \square w_2) \\ &= f_1(a \otimes b)a(bw_2 \square w_1) + f_1(b \otimes a)b(w_2 \square aw_1) = v \square u. \end{aligned}$$

case $a = b$ and $f_1(a \otimes a) = f_2(a \otimes a)$.

$$\begin{aligned} u \square v &= f_1(a \otimes a)a(w_1 \square aw_2) + f_1(a \otimes a)a(aw_1 \square w_2) \\ &= f_1(a \otimes a)a(aw_2 \square w_1) + f_1(a \otimes a)a(w_2 \square aw_1) = v \square u. \end{aligned}$$

case $a = b$ and $f_2(a \otimes a) = 1 - f_1(a \otimes a)$. There exist two words w_3 and w_4 , not necessarily non-empty, not starting by a and two positive integers k and l such that $w_1 = \underbrace{a \dots a}_{k \text{ times}} w_3$ and $w_2 = \underbrace{a \dots a}_{l \text{ times}} w_4$. First of all, by induction,

$$\underbrace{a \dots a}_{k \text{ times}} \square \underbrace{a \dots a}_{l \text{ times}} = \underbrace{a \dots a}_{k+l \text{ times}} .$$

Besides, relations satisfied by \square enjoin $f_1(a \otimes c) = 1$ and $f_2(c \otimes a) = 0$ for any letter $c \neq a$. So,

$$u \square v = (\underbrace{a \dots a}_{k \text{ times}} \square \underbrace{a \dots a}_{l \text{ times}})(w_3 \square w_4) = (\underbrace{a \dots a}_{l \text{ times}} \square \underbrace{a \dots a}_{k \text{ times}})(w_4 \square w_3) = v \square u.$$

As a consequence, \square is a commutative product.

Let now w_1, w_2 and w_3 be three words such that $\text{length}(w_1) + \text{length}(w_2) + \text{length}(w_3) = n + 1$. Then there exist three letters a, b and c and three words w_4, w_5 and w_6 (not necessarily non-empty) such that $w_1 = aw_4, w_2 = bw_5$ and $w_3 = cw_6$. Then, by induction, we get:

case a, b and c distinct.

$$(w_1 \square w_2) \square w_3 = f_1(a \otimes b)f_1(a \otimes c)a[(w_4 \square bw_5) \square cw_6] + f_1(a \otimes b)f_1(c \otimes a)c[a(w_4 \square bw_5) \square w_6] \\ + f_1(b \otimes a)f_1(b \otimes c)b[(aw_4 \square w_5) \square cw_6] + f_1(b \otimes a)f_1(c \otimes b)c[b(aw_4 \square w_5) \square w_6]$$

and

$$w_1 \square (w_2 \square w_3) = f_1(b \otimes c)f_1(a \otimes b)a[w_4 \square b(w_5 \square cw_6)] + f_1(b \otimes c)f_1(b \otimes a)b[aw_4 \square (w_5 \square cw_6)] \\ + f_1(c \otimes b)f_1(a \otimes c)a[w_4 \square c(bw_5 \square w_6)] + f_1(c \otimes b)f_1(c \otimes a)c[aw_4 \square (bw_5 \square w_6)].$$

However

$$(w_4 \square bw_5) \square cw_6 = w_4 \square (bw_5 \square cw_6) = f_1(b \otimes c)w_4 \square b(w_5 \square cw_6) + f_1(c \otimes b)w_4 \square c(bw_5 \square w_6), \\ aw_4 \square (bw_5 \square w_6) = (aw_4 \square bw_5) \square w_6 = f_1(a \otimes b)a(w_4 \square bw_5) \square w_6 + f_1(b \otimes a)b(aw_4 \square w_5) \square w_6, \\ \text{and } f_1 \text{ satisfies } f_1(x \otimes y)f_1(y \otimes z)(f_1(x \otimes z) - 1) = 0 \text{ for any set } \{x, y, z\} \in X. \text{ Thus,} \\ (w_1 \square w_2) \square w_3 = w_1 \square (w_2 \square w_3).$$

case $a = b$ and $(a \neq c)$. By commutativity it is the same case as $(a = c$ and $b \neq a)$ or $(b = c$ and $a \neq b)$.

$$(w_1 \square w_2) \square w_3 = f_1(a \otimes a)f_1(a \otimes c)a[(w_4 \square aw_5) \square cw_6] + f_1(a \otimes a)f_1(c \otimes a)c[a(w_4 \square aw_5) \square w_6] \\ + f_2(a \otimes a)f_1(a \otimes c)a[(aw_4 \square w_5) \square cw_6] + f_2(a \otimes a)f_1(c \otimes a)c[a(aw_4 \square w_5) \square w_6]$$

and

$$w_1 \square (w_2 \square w_3) = f_1(a \otimes c)f_1(a \otimes a)a[w_4 \square a(w_5 \square cw_6)] + f_1(a \otimes c)f_2(a \otimes a)a[aw_4 \square (w_5 \square cw_6)] \\ + f_1(c \otimes a)f_1(a \otimes c)a[w_4 \square c(aw_5 \square w_6)] + f_1(c \otimes a)^2c[aw_4 \square (aw_5 \square w_6)].$$

However

$$(w_4 \square aw_5) \square cw_6 = w_4 \square (aw_5 \square cw_6) = f_1(a \otimes c)w_4 \square a(w_5 \square cw_6) + f_1(c \otimes a)w_4 \square c(aw_5 \square w_6), \\ aw_4 \square (aw_5 \square w_6) = (aw_4 \square aw_5) \square w_6 = f_1(a \otimes a)a(w_4 \square aw_5) \square w_6 + f_2(a \otimes a)a(aw_4 \square w_5) \square w_6, \\ \text{and } f_1 \text{ satisfies}$$

1. If $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$ then

- (a) $f_1(a \otimes b)f_1(b \otimes a)[f_1(a \otimes a) - 1] = 0$,
- (b) $f_1(a \otimes a)f_1(a \otimes b)[f_1(a \otimes b) - 1] = 0$,
- (c) $f_1(a \otimes a)f_1(b \otimes a)[f_1(b \otimes a) - 1] = 0$.

2. If $f_1(a \otimes a) = 1 - f_2(a \otimes a)$ then $f_1(a \otimes c) = 1$ and $f_1(c \otimes a) = 0$.

Thus, $(w_1 \square w_2) \square w_3 = w_1 \square (w_2 \square w_3)$.

case $a = b = c$ and $f_1(a \otimes a) = f_2(a \otimes a)$.

$$(w_1 \square w_2) \square w_3 = f_1(a \otimes a)^2 a [(w_4 \square a w_5) \square a w_6] + f_1(a \otimes a)^2 a [a (w_4 \square a w_5) \square w_6] \\ + f_1(a \otimes a)^2 a [(a w_4 \square w_5) \square a w_6] + f_1(a \otimes a)^2 a [a (a w_4 \square w_5) \square w_6]$$

and

$$w_1 \square (w_2 \square w_3) = f_1(a \otimes a)^2 a [w_4 \square a (w_5 \square a w_6)] + f_1(a \otimes a)^2 a [a w_4 \square (w_5 \square a w_6)] \\ + f_1(a \otimes a)^2 a [w_4 \square a (a w_5 \square w_6)] + f_1(a \otimes a)^2 a [a w_4 \square (a w_5 \square w_6)].$$

Thus, $(w_1 \square w_2) \square w_3 = w_1 \square (w_2 \square w_3)$.

case $a = b = c$ and $f_2(a \otimes a) = 1 - f_1(a \otimes a)$. There exist three words w_7 , w_8 and w_9 not necessarily non-empty, not starting by a and three positive integers k , l and m such that $w_1 = \underbrace{a \dots a}_{k \text{ times}} w_7$, $w_2 = \underbrace{a \dots a}_{l \text{ times}} w_8$ and $w_3 = \underbrace{a \dots a}_{m \text{ times}} w_9$. Besides, relations satisfied by \square enjoin $f_1(a \otimes c) = 1$ and $f_2(c \otimes a) = 0$ for any letter $c \neq a$. So,

$$(w_1 \square w_2) \square w_3 = \left[\underbrace{(a \dots a)}_{k \text{ times}} \square \underbrace{(a \dots a)}_{l \text{ times}} \square \underbrace{(a \dots a)}_{k \text{ times}} \right] \left[(w_7 \square w_8) \square w_9 \right] \\ = \underbrace{a \dots a}_{k+l+m \text{ times}} \left[(w_7 \square w_8) \square w_9 \right] \\ = \left[\underbrace{a \dots a}_{k \text{ times}} \square \left(\underbrace{a \dots a}_{l \text{ times}} \square \underbrace{a \dots a}_{k \text{ times}} \right) \right] \left[w_7 \square (w_8 \square w_9) \right] = w_1 \square (w_2 \square w_3).$$

□

Corollary 11. Let \mathbb{K} be a field of characteristic 0, let X be a countable alphabet and let \square be a weak shuffle product on $\mathbb{K}\langle X \rangle$.

1. There exists at most one letter a such that $f_1(a \otimes a) = 1 - f_2(a \otimes a)$.
2. If there exists a letter a such that $f_1(a \otimes a) = 1 - f_2(a \otimes a)$ then, for any word u and v , the calculation of $u \square v$ does not depend on the value of $f_1(a \otimes a)$.
3. If $f_1(a \otimes b) = f_1(b \otimes a) = 1$ then $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = 1$, $f_1(a \otimes c) = f_1(b \otimes c) \in \{0, 1\}$ and $f_1(c \otimes a) = f_1(c \otimes b) \in \{0, 1\}$ for any $c \in X \setminus \{a, b\}$.

Proof. 1. If there are two letters a and b such that $a \neq b$, $f_1(a \otimes a) = 1 - f_2(a \otimes a)$ and $f_1(b \otimes b) = 1 - f_2(b \otimes b)$ then $1 = f_1(a \otimes b) = 0$ and $0 = f_1(b \otimes a) = 1$. Contradiction.

2. Let a such that $f_1(a \otimes a) = 1 - f_2(a \otimes a)$. If u and v are words in $X^* \setminus aX^*$, since $f_1(a \otimes b) = 1$ and $f_1(b \otimes a) = 0$ for any $b \neq a$, there does not exist any triple (w, u', v') such that $u \square v = w(a u' \square a v')$.

3. If $f_1(a \otimes b) = f_1(b \otimes a) = 1$ then, the fact that $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = 1$ comes directly from relations 2 given in Theorem 10. To prove $f_1(a \otimes c) = f_1(b \otimes c) \in \{0, 1\}$ and $f_1(c \otimes a) = f_1(c \otimes b) \in \{0, 1\}$ for any $c \in X \setminus \{a, b\}$, we use the relation $f_1(x \otimes y)f_1(y \otimes z)[f_1(x \otimes z) - 1] = 0$ for any $x, y, z \in X$. □

Proposition 12. *Let \mathbb{K} be a field of characteristic 0, X be a countable alphabet and \square a weak shuffle product on $\mathbb{K}\langle X \rangle$. We denote by T the set $T = \{a \in X, f_1(a \otimes a) \in \mathbb{K} \setminus \{0, 1\}\}$. We assume $T \neq \emptyset$; so T is a singleton $\{a\}$. Let \square' be the weak shuffle product defined by*

- $f'_1(u \otimes v) = f_1(u \otimes u)$ for any $u \otimes v \in X \otimes X \setminus \{a \otimes a\}$,
- $f'_1(a \otimes a) = 1$ and $f'_2(a \otimes a) = 1$.

Then, there exists an algebra isomorphism between $(\mathbb{K}\langle X \rangle, \square)$ and $(\mathbb{K}\langle X \rangle, \square')$.

Proof. Thanks to Corollary 11, we know that the weak shuffle \square does not depend on the value of $f_1(a \otimes a)$. We define $\psi : (\mathbb{K}\langle X \rangle, \square) \rightarrow (\mathbb{K}\langle X \rangle, \square')$ by:

$$\psi(w) = \begin{cases} w & \text{if } w \notin aX^*, \\ \frac{1}{n!}w & \text{if } w = \underbrace{a \dots a}_n w_1 \text{ with } w_1 \notin aX^*. \end{cases}$$

Since $f_1(a \otimes b) = 1$ and $f_1(b \otimes a) = 0$ for any $b \in X \setminus \{a\}$, the linear map ψ is an algebra morphism. It is trivially an isomorphism. □

Proposition 13. *Let \mathbb{K} be a field of characteristic 0, let X be an alphabet of cardinality 2 or 3 and let \square be a weak shuffle product on $\mathbb{K}\langle X \rangle$. Let \square' be the weak shuffle product defined by*

- $f'_1(a \otimes b) = 1$ and $f'_1(b \otimes a) = 0$ for any $(a \otimes b) \in X \otimes X$ such that $a \neq b$ and $f_1(a \otimes b) \notin \{0, 1\}$.
- $f'_1(a \otimes b) = f_1(a \otimes b)$ for any $(a \otimes b) \in X \otimes X$ such that $a \neq b$ and $f_1(a \otimes b) \in \{0, 1\}$.
- $f'_i(a \otimes a) = f_i(a \otimes a)$ for any $a \in X$ and any $i \in \{1, 2\}$.

Then, there exists an algebra isomorphism between $(\mathbb{K}\langle X \rangle, \square)$ and $(\mathbb{K}\langle X \rangle, \square')$.

Proof. If $X = \{a, b\}$ then there is an one-parameter family of weak shuffle products \square such that $f_1(a \otimes b) \notin \{0, 1\}$. They are defined by $f_1(a \otimes b) = k \in \mathbb{K} \setminus \{0, 1\}$ and $f_1(b \otimes a) = f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = 0$. We define \square' by changing k in 1. The map φ defined by

$$\varphi(w) = \begin{cases} \frac{1}{k^n}w & \text{if } w = \underbrace{a \dots a}_n w' \text{ with } w' \in bX^*, \\ w & \text{else,} \end{cases}$$

is an algebra isomorphism between $(\mathbb{K}\langle X \rangle, \square)$ and $(\mathbb{K}\langle X \rangle, \square')$

Let us now consider the case $X = \{a, b, c\}$. Without loss of generality we assume $f_1(a \otimes b) = k \in \mathbb{K} \setminus \{0, 1\}$. The characterisation of weak shuffle products given in Theorem 10 leads to the following relations:

- $f_1(b \otimes a) = f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = 0,$
- $f_1(a \otimes c)f_1(c \otimes a) = 0,$
- $f_1(b \otimes c)f_1(c \otimes b) = 0,$
- $f_1(a \otimes c)f_1(c \otimes b) = 0,$
- $f_1(b \otimes c)f_1(c \otimes a) = 0,$

- $f_1(u \otimes v)f_1(v \otimes w)[f_1(u \otimes w) - 1] = 0$ where $\{u, v, w\} = X$.

Thus, the weak shuffle product \square is one of the following:

1. $f_1(a \otimes c) = f_1(b \otimes c) = f_1(c \otimes a) = f_1(c \otimes b) = 0$ and $f_1(c \otimes c) = f_2(c \otimes c) \in \{0, 1\}$.
2. $f_1(a \otimes c) = 1$, $f_1(b \otimes c) = p \in \mathbb{K}^*$ and $f_1(c \otimes a) = f_1(c \otimes b) = f_1(c \otimes c) = f_2(c \otimes c) = 0$,
3. $f_1(a \otimes c) = 1$, $f_1(b \otimes c) = 1$, $f_1(c \otimes a) = f_1(c \otimes b) = 0$ and $f_1(c \otimes c) = f_2(c \otimes c) = 1$,
4. $f_1(a \otimes c) = f_1(b \otimes c) = 0$, $f_1(c \otimes a) = p \in \mathbb{K}^*$, $f_1(c \otimes b) = 1$ and $f_1(c \otimes c) = f_2(c \otimes c) = 0$,
5. $f_1(a \otimes c) = f_1(b \otimes c) = 0$, $f_1(c \otimes a) = 1$, $f_1(c \otimes b) = 1$ and $f_1(c \otimes c) = f_2(c \otimes c) = 1$,
6. $f_1(a \otimes c) = f_1(b \otimes c) = 0$, $f_1(c \otimes a) = 1$, $f_1(c \otimes b) = 1$ and $f_1(c \otimes c) = 1 - f_2(c \otimes c)$,
7. $f_1(a \otimes c) = f_1(b \otimes c) = f_1(c \otimes a) = 0$, $f_1(c \otimes b) = p \in \mathbb{K}^*$ and $f_1(c \otimes c) = f_2(c \otimes c) = 0$,
8. $f_1(a \otimes c) = f_1(b \otimes c) = f_1(c \otimes a) = 0$, $f_1(c \otimes b) = 1$ and $f_1(c \otimes c) = f_2(c \otimes c) = 1$,
9. $f_1(a \otimes c) = p \in \mathbb{K}^*$, $f_1(b \otimes c) = f_1(c \otimes a) = f_1(c \otimes b) = 0$ and $f_1(c \otimes c) = f_2(c \otimes c) = 0$,
10. $f_1(a \otimes c) = 1$, $f_1(b \otimes c) = f_1(c \otimes a) = f_1(c \otimes b) = 0$ and $f_1(c \otimes c) = f_2(c \otimes c) = 1$,

We define \square' by $f_1'(a \otimes b) = 1$ and $f_1'(u \otimes v) = f_1(u \otimes v)$ if $u \otimes v \neq a \otimes b$. Let φ_1 and φ_2 be the maps defined by: for any word w ,

$$\varphi_1(w) = \begin{cases} \frac{1}{k^n} w & \text{if } w = \underbrace{a \dots a}_n w' \text{ with } w' \in bX^*, \\ w & \text{else,} \end{cases}$$

and

$$\varphi_2(w) = \begin{cases} \frac{1}{k^{n_1 + \dots + n_s}} w & \text{if } w = \underbrace{c \dots c}_{q_1 \text{ times}} \underbrace{a \dots a}_{n_1 \text{ times}} \underbrace{c \dots c}_{q_2 \text{ times}} \dots \underbrace{c \dots c}_{q_s \text{ times}} \underbrace{a \dots a}_{n_s \text{ times}} \underbrace{c \dots c}_{q_{s+1} \text{ times}} w' \text{ with } w' \in bX^* \\ & \text{and } (q_1, \dots, q_{s+1}) \in \mathbb{N}^{s+1}, \\ w & \text{else.} \end{cases}$$

From case 1 to case 3 and from case 9 to case 10 the map φ_1 is an algebra isomorphism between $(\mathbb{K}\langle X \rangle, \square)$ and $(\mathbb{K}\langle X \rangle, \square')$. From case 4 to case 8 the map φ_2 is an algebra isomorphism between $(\mathbb{K}\langle X \rangle, \square)$ and $(\mathbb{K}\langle X \rangle, \square')$.

If maps f_1' and f_2' do not take their values in $\{0, 1\}$ we apply the previous process once again to \square' . And then, we find a weak shuffle product \square'' such that $f_1''(u \otimes v), f_2''(u \otimes v) \in \{0, 1\}$ for any $(u \otimes v) \in X \otimes X$. \square

Conjecture 14. *Proposition 13 is still true for any countable alphabet.*

Remark. If X is an alphabet such that $\{a, b, c, d\} \subset X$ and $f_1(a \otimes b) \notin \{0, 1\}$ then relations

1. $f_1(a \otimes x)f_1(x \otimes a) = 0$,
2. $f_1(b \otimes x)f_1(x \otimes b) = 0$,
3. $f_1(a \otimes x)f_1(x \otimes b) = 0$,
4. $f_1(b \otimes x)f_1(x \otimes a) = 0$,

are still satisfied for any letter $x \in X$. However, if $x, y \in X \setminus \{a, b\}$, even if they satisfy relations given in Theorem 10, it is hard to anticipate the part of x facing y .

2.2 Weak shuffle products on $\mathbb{K}\langle\{a, b\}\rangle$

Let $X = \{a, b\}$ be an alphabet of cardinality 2. By using the characterisation given in Theorem 10, there are 10 families of weak shuffle products defined on $\mathbb{K}\langle X \rangle$. Let C be the 6-tuple $C = (f_1(a \otimes b), f_1(b \otimes a), f_1(a \otimes a), f_2(a \otimes a), f_1(b \otimes b), f_2(b \otimes b))$. If $k \in \mathbb{K}^*$ and $\alpha \in \mathbb{K}$ then C is one of the following 6-tuples

$$\begin{aligned} C_1 &= (0, 0, 0, 0, 0, 0), & C_2 &= (k, 0, 0, 0, 0, 0), & C_3 &= (1, 0, 1, 1, 0, 0), \\ C_4 &= (1, 0, 0, 0, 1, 1), & C_5 &= (0, 0, 1, 1, 0, 0), & C_6 &= (0, 0, 1, 1, 1, 1), \\ C_7 &= (1, 0, \alpha, 1 - \alpha, 0, 0), & C_8 &= (1, 0, \alpha, 1 - \alpha, 1, 1), & C_9 &= (1, 0, 1, 1, 1, 1), \\ C_{10} &= (1, 1, 1, 1, 1, 1). \end{aligned}$$

For any $n \in \llbracket 1, 10 \rrbracket$, we denote by \square_n the weak shuffle product associated to C_n . The concatenation of two words u and v is denoted by uv . The empty word is denoted by 1.

Case $n = 2$. Thanks to Proposition 13, for any $k \in \mathbb{K}^*$ the weak shuffle product defined by C_2 is isomorphic to the case $(1, 0, 0, 0, 0, 0)$. Let u and v be two non-empty words. Then

$$u \square_2 v = \begin{cases} k^n uv & \text{if } (u = \underbrace{a \dots a}_{n \text{ times}} \text{ and } v = bw \text{ with } w \in X^*) \\ & \text{or } (v = a \dots a \text{ and } u = bw \text{ with } w \in X^*), \\ 0 & \text{else.} \end{cases}$$

Cases $n = 3$ and $n = 7$. Thanks to Proposition 12 the weak shuffle products defines by C_3 and C_7 are isomorphic. Let u and v be two non-empty words. Then

$$u \square_3 v = \begin{cases} uv & \text{if } (u = a \dots a \text{ and } v = bw \text{ with } w \in X^*) \\ & \text{or } (v = a \dots a \text{ and } u = bw \text{ with } w \in X^*), \\ \binom{k+l}{k} \underbrace{a \dots a}_{k+l \text{ times}} w & \text{if } (u = \underbrace{a \dots a}_{k \text{ times}} \text{ and } v = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^* \cup \{1\}) \\ & \text{or } (v = \underbrace{a \dots a}_{k \text{ times}} \text{ and } u = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^* \cup \{1\}), \\ 0 & \text{else,} \end{cases}$$

$$\text{and } u \square_7 v = \begin{cases} uv & \text{if } (u = a \dots a \text{ and } v = bw \text{ with } w \in X^*) \\ & \text{or } (v = a \dots a \text{ and } u = bw \text{ with } w \in X^*), \\ \underbrace{a \dots a}_{k+l \text{ times}} w & \text{if } (u = \underbrace{a \dots a}_{k \text{ times}} \text{ and } v = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^* \cup \{1\}) \\ & \text{or } (v = \underbrace{a \dots a}_{k \text{ times}} \text{ and } u = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^* \cup \{1\}), \\ 0 & \text{else.} \end{cases}$$

Case $n = 5$. Let u and v be two non-empty words. Then

$$u \square_5 v = \begin{cases} \binom{k+l-1}{k} \underbrace{a \dots a}_{k+l \text{ times}} w & \text{if } (u = \underbrace{a \dots a}_{k \text{ times}} \text{ and } v = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^*) \\ & \text{or } (v = \underbrace{a \dots a}_{k \text{ times}} \text{ and } u = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^*), \\ \binom{k+l}{k} \underbrace{a \dots a}_{k+l \text{ times}} & \text{if } u = \underbrace{a \dots a}_{k \text{ times}} \text{ and } v = \underbrace{a \dots a}_{l \text{ times}}, \\ 0 & \text{else.} \end{cases}$$

Case $n = 6$. Let u and v be two non-empty words. Then

$$u \square_6 v = \begin{cases} \binom{k+l-1}{k} \underbrace{a \dots a}_{k+l \text{ times}} w & \text{if } (u = \underbrace{a \dots a}_{k \text{ times}} \text{ and } v = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^*) \\ & \text{or } (v = \underbrace{a \dots a}_{k \text{ times}} \text{ and } u = \underbrace{a \dots a}_{l \text{ times}} w \text{ with } w \in bX^*), \\ \binom{k+l}{k} \underbrace{a \dots a}_{k+l \text{ times}} & \text{if } u = \underbrace{a \dots a}_{k \text{ times}} \text{ and } v = \underbrace{a \dots a}_{l \text{ times}}, \\ \binom{k+l-1}{k} \underbrace{b \dots b}_{k+l \text{ times}} w & \text{if } (u = \underbrace{b \dots b}_{k \text{ times}} \text{ and } v = \underbrace{b \dots b}_{l \text{ times}} w \text{ with } w \in aX^*) \\ & \text{or } (v = \underbrace{b \dots b}_{k \text{ times}} \text{ and } u = \underbrace{b \dots b}_{l \text{ times}} w \text{ with } w \in aX^*), \\ \binom{k+l}{k} \underbrace{b \dots b}_{k+l \text{ times}} & \text{if } u = \underbrace{b \dots b}_{k \text{ times}} \text{ and } v = \underbrace{b \dots b}_{l \text{ times}}, \\ 0 & \text{else.} \end{cases}$$

Case $n = 4$. First, it is natural to ask whether or not this case is isomorphic to the case with $n = 3$? In fact, not. A counter-example is given by the elements u of degree 2 such that $u^2 = 0$. Indeed,

1. with the case $n = 4$, if $u = \lambda aa + \mu bb + \sigma ab + \tau ba$ then

$$\begin{aligned} u^2 = & 6\mu^2 bbbb + 2\tau^2 baba + 2\lambda\mu aabb + 2\lambda\tau aaba + 6\mu\sigma abbb \\ & + 2\mu\tau (babb + bbab + bbba) + 2\sigma\tau (abab + abba). \end{aligned}$$

$$\text{So } u^2 = 0 \iff \mu = \tau = 0 \text{ and } \left\{ u \in \mathbb{K}\langle\{a, b\}^*\rangle, \text{length}(u) = 2 \text{ and } u^2 = 0 \right\} = \text{Span}(aa, ab).$$

2. with the case $n = 3$, if $u = \lambda aa + \mu bb + \sigma ab + \tau ba$ then

$$u^2 = 6\lambda^2 aaaa + 2\lambda\mu aabb + 6\lambda\sigma aaab + 2\lambda\tau aaba.$$

$$\text{So } u^2 = 0 \iff \lambda = 0 \text{ and } \left\{ u \in \mathbb{K}\langle\{a, b\}^*\rangle, \text{length}(u) = 2 \text{ and } u^2 = 0 \right\} = \text{Span}(bb, ab, ba).$$

Let u and v be two non-empty words. Then

1. If $u = \underbrace{a \dots a}_m u'$ and $u', v \in bX^* \cup \{1\}$ then

$$u \square_4 v = v \square_4 u = \underbrace{a \dots a}_m (u' \square_4 v).$$

2. If $u = \underbrace{b \dots b}_{m_1} u', v = \underbrace{b \dots b}_{m_2} v'$ and $u', v' \in aX^* \cup \{1\}$ then

$$\begin{aligned} u \square_4 v &= \sum_{k=0}^{m_2-1} \binom{m_1+k-1}{k} \underbrace{b \dots b}_{m_1+k \text{ times}} (u' \square_4 \underbrace{b \dots b}_{m_2-k \text{ times}} v') \\ &+ \sum_{k=0}^{m_1-1} \binom{m_2+k-1}{k} \underbrace{b \dots b}_{m_2+k \text{ times}} (\underbrace{b \dots b}_{m_1-k \text{ times}} u' \square_4 v') \\ &= v \square_4 u \end{aligned}$$

3. If $u, v \in aX^*$ then $u \square_4 v = v \square_4 u = 0$.

Cases $n = 8$ and $n = 9$. We recall that the case $n = 8$ does not depend on $\alpha \in \mathbb{K}$. Thanks to Proposition 12 the weak shuffle products defined by C_8 and C_9 are isomorphic. Let u and v be two non-empty words. Then

1. If $u = \underbrace{a \dots a}_m u'$ and $u', v \in bX^* \cup \{1\}$ then

$$u \square_9 v = v \square_9 u = \underbrace{a \dots a}_m (u' \square_9 v) = u \square_8 v = v \square_8 u.$$

2. If $u = \underbrace{b \dots b}_{m_1} u', v = \underbrace{b \dots b}_{m_2} v'$ and $u', v' \in aX^* \cup \{1\}$ then

$$\begin{aligned} u \square_9 v &= \sum_{k=0}^{m_2-1} \binom{m_1+k-1}{k} \underbrace{b \dots b}_{m_1+k} (u' \square_9 \underbrace{b \dots b}_{m_2-k} v') \\ &\quad + \sum_{k=0}^{m_1-1} \binom{m_2+k-1}{k} \underbrace{b \dots b}_{m_2+k} (\underbrace{b \dots b}_{m_1-k} u' \square_9 v') \\ &= v \square_9 u = u \square_8 v = v \square_8 u. \end{aligned}$$

3. If $u = \underbrace{a \dots a}_k u', v = \underbrace{a \dots a}_l v'$ and $u', v' \in bX^* \cup \{1\}$ then

$$u \square_9 v = v \square_9 u = \binom{k+l}{k} \underbrace{a \dots a}_{k+l} (u' \square_9 v'),$$

and

$$u \square_8 v = v \square_8 u = \underbrace{a \dots a}_{k+l} (u' \square_8 v').$$

From the previous calculations, we have the following consequence:

Corollary 15. *Let v and w be two words. Then $v \square_9 w \neq 0$.*

Remark. For cases $n \in \{4, 8, 9\}$, since $f_1(a \otimes b) = 1$ and $f_1(b \otimes a) = 0$, the calculation of $u \square_n v$ where $u = \underbrace{b \dots b}_{m_1} u', v = \underbrace{b \dots b}_{m_2} v'$ and $u', v' \in aX^* \cup \{1\}$ does not depend on the values of $f_1(a \otimes a)$ nor $f_2(a \otimes a)$. We give the value of $u \square_4 v (= u \square_8 v = u \square_9 v)$ for some example couple $(u, v) \in (bX^*)^2$. For some examples of pairs $(x, p) \in X \times \mathbb{N}^*$, to lighten the notation, we write x^p instead of $\underbrace{x \dots x}_p$.

Let (m, s, p, r) be a quadruple of positive integers. Then:

$$b^m a^s \square_4 b^p a^r = \sum_{k=0}^{p-1} \binom{m+k-1}{k} b^{m+k} a^s b^{p-k} a^r + \sum_{k=0}^{m-1} \binom{p+k-1}{k} b^{p+k} a^r b^{m-k} a^s.$$

Let (m, s, p, r, t) be a quintuple of positive integers such that $m \geq 2$. Then:

$$\begin{aligned} b^m a^s \square_4 b^p a^r b^t &= \sum_{k=0}^{p-1} \binom{m+k-1}{k} b^{m+k} a^s b^{p-k} a^r b^t + \sum_{k=0}^t \binom{m+k-1}{k} b^p a^r b^{m+k} a^s b^{t-k} \\ &\quad + \sum_{\substack{f+g=m \\ f \in \mathbb{N}^* \\ g \in \mathbb{N}^*}} \sum_{k=0}^t \binom{f+p-1}{f} \binom{g+k-1}{k} b^{p+f} a^r b^{g+k} a^s b^{t-k}. \end{aligned}$$

Proposition 16. Let \square_9 be the weak shuffle product defined by C_9 . Let p be a positive integer and $n \in \{1, 2, 3\}$. We denote by $K_{(n,p)}$ the set

$$K_{(n,p)} = \left\{ u = \sum_{\substack{w \in X^* \\ \text{length}(w)=n}} \lambda_w w, u^p = 0 \right\}.$$

Then, $K_{(n,p)} = \{0\}$.

Proof. We equip X^* with the lexicographic order. For any words v and w we denote by $\max(v \square w)$ the greatest word of length $l = \text{length}(v) + \text{length}(w)$ which appears in $v \square w$ for the lexicographic order.

If $u = \sum_{\substack{w \in X^* \\ \text{length}(w)=n}} \lambda_w w$ then

$$u^p = \sum_{\substack{w \in X^* \\ \text{length}(w)=n}} \lambda_w^p (w \square_9 \dots \square_9 w) + \sum_{l=2}^{\min(p,n)} \sum_{(\alpha_1, \dots, \alpha_l) \models p} \sum_{\substack{w_1 < \dots < w_l \in X^* \\ \forall i \text{ length}(w_i)=n}} \lambda_{w_1}^{\alpha_1} \dots \lambda_{w_l}^{\alpha_l} (w_1 \square_9 \dots \square_9 w_l).$$

1. If $n = 1$ then the result is trivial.

2. If $n = 2$ then

$$aa^p = \frac{(2p)!}{2^p} \underbrace{a \dots a}_{2p \text{ times}}, \quad ab^p = (p!)^2 \underbrace{a \dots a}_{p \text{ times}} \underbrace{b \dots b}_{p \text{ times}}, \quad ba^p = p! \underbrace{ba \dots ba}_{p \text{ times}}, \quad bb^p = \frac{(2p)!}{2^p} \underbrace{b \dots b}_{2p \text{ times}},$$

and

$$\max(aa^k \square_9 ab^l \square_9 ba^m \square_9 bb^n) = \underbrace{a \dots a}_{2k+l \text{ times}} \underbrace{b \dots b}_{2n+l \text{ times}} \underbrace{ba \dots ba}_{m \text{ times}}.$$

Thus $\lambda_{aa} = \lambda_{bb} = \lambda_{ba} = \lambda_{ba} = 0$.

3. If $n = 3$ then

$$\begin{aligned} w_1 = aaa^p &= \frac{(3p)!}{(3!)^p} \underbrace{a \dots a}_{3p \text{ times}}, & w_2 = aab^p &= \frac{(2p)!p!}{2^p} \underbrace{a \dots a}_{2p \text{ times}} \underbrace{b \dots b}_{p \text{ times}}, \\ w_3 = aba^p &= (p!)^2 \underbrace{a \dots a}_{p \text{ times}} \underbrace{ba \dots ba}_{p \text{ times}}, & w_4 = abb^p &= \frac{(2p)!p!}{2^p} \underbrace{a \dots a}_{p \text{ times}} \underbrace{b \dots b}_{2p \text{ times}}, \\ w_5 = baa^p &= p! \underbrace{baa \dots baa}_{p \text{ times}}, & w_6 = bbb^p &= \frac{(3p)!}{(3!)^p} \underbrace{b \dots b}_{3p \text{ times}}. \end{aligned}$$

For bab^p and bba^p , there are several terms in the result. For bab^p we will use $w_7 = \underbrace{bab \dots bab}_{p \text{ times}}$

and, for bba^p we will use $w_8 = \underbrace{b \dots b}_{p \text{ times}} \underbrace{ba \dots ba}_{p \text{ times}}$. In fact, for the lexicographic order, we use

the maximal term obtained in each product. For any i determine how build w_i by doing the weak shuffle of p words of length 3. We get $\lambda_{aaa} = \lambda_{bbb} = \lambda_{aba} = \lambda_{baa} = \lambda_{aab} = \lambda_{abb} = \lambda_{bab} = \lambda_{bba} = 0$.

□

Conjecture 17. Let \square_9 be the weak shuffle product defined by C_9 . For any positive integers p and n , we have $K_{(n,p)} = \{0\}$.

Remarks.

1. By induction we can express $\max(u \square_9 v)$ for any words u and v .

Case w_1 and w_2 are in aX^* . There exist $\alpha, \beta \in \mathbb{N}^*$ and $w'_1, w'_2 \in bX^* \cup \{1\}$ such that $w_1 = \underbrace{a \dots a}_{\alpha \text{ times}} w'_1$ and $w_2 = \underbrace{a \dots a}_{\beta \text{ times}} w'_2$. Then,

$$\max(w_1 \square_9 w_2) = \underbrace{a \dots a}_{\alpha + \beta \text{ times}} \max(w'_1 \square_9 w'_2).$$

Case $w_1 \in aX^*$ and $w_2 \in bX^*$. There exist $\alpha \in \mathbb{N}^*$ and $w'_1 \in bX^* \cup \{1\}$ such that $w_1 = \underbrace{a \dots a}_{\alpha \text{ times}} w'_1$. Then,

$$\max(w_1 \square_9 w_2) = \underbrace{a \dots a}_{\alpha \text{ times}} \max(w'_1 \square_9 w_2).$$

Case w_1 and w_2 are in bX^* . There exist $\alpha, \beta \in \mathbb{N}^*$, $p, q \in \mathbb{N}$ (they are not necessarily different from 0) and $w'_1, w'_2 \in bX^* \cup \{1\}$ such that $w_1 = \underbrace{b \dots b}_{\alpha \text{ times}} \underbrace{a \dots a}_{p \text{ times}} w'_1$ and $w_2 =$

$\underbrace{b \dots b}_{\beta \text{ times}} \underbrace{a \dots a}_{q \text{ times}} w'_2$. Thus,

- If $0 < q < p$ then

$$\max(w_1 \square_9 w_2) = \underbrace{b \dots b}_{\alpha + \beta - 1 \text{ times}} \underbrace{a \dots a}_{q \text{ times}} \max(\underbrace{b a \dots a}_{p \text{ times}} w'_1 \square_9 w'_2).$$

- If $0 < p < q$ then

$$\max(w_1 \square_9 w_2) = \underbrace{b \dots b}_{\alpha + \beta - 1 \text{ times}} \underbrace{a \dots a}_{p \text{ times}} \max(w'_1 \square_9 \underbrace{b a \dots a}_{q \text{ times}} w'_2).$$

- If $0 < p$ and $p = q$ then $\max(w_1 \square_9 w_2) = \max(\tilde{w}_1, \tilde{w}_2)$ where

$$\tilde{w}_1 = \underbrace{b \dots b}_{\alpha + \beta - 1 \text{ times}} \underbrace{a \dots a}_{q \text{ times}} \max(\underbrace{b a \dots a}_{p \text{ times}} w'_1 \square_9 w'_2)$$

and

$$\tilde{w}_2 = \underbrace{b \dots b}_{\alpha + \beta - 1 \text{ times}} \underbrace{a \dots a}_{p \text{ times}} \max(w'_1 \square_9 \underbrace{b a \dots a}_{q \text{ times}} w'_2).$$

- If $p = 0$ (respectively $q = 0$) then $w_1 = \underbrace{b \dots b}_{\alpha \text{ times}}$ (respectively $w_2 = \underbrace{b \dots b}_{\beta \text{ times}}$) and

$$\max((w_1 \square_9 w_2)) = w_1 w_2 \text{ (respectively } \max((w_1 \square_9 w_2)) = w_2 w_1).$$

For instance,

$$\max(ab \square_9 abaa) = aa \max(b \square_9 baa) = abbbaa,$$

$$\max(bba \square_9 baa) = bbabaa,$$

$$\max(bbbaaabba \square_9 bbaabbba) = bbbbaa \max(baaabba \square_9 bbba) = bbbbaabbabaaabba.$$

2. For $p = 2$ Conjecture 17 is implied by the statement "Let n be a positive integer, let w_1 , w_2 and w be three non-empty words of length n such that $w_1 \leq w_2 \leq w$ and $w_1 < w$. Then $\max(w_1 \square_9 w_2) < \max(w \square_9 w)$." We attend a reasoning by induction but there are some obstructions. Indeed, it leads us to compare $\max(u_1 \square_9 u_2)$ and $\max(u_3 \square_9 u_4)$ where $u_1 \leq u_3$, $u_2 \leq u_4$, $\text{length}(u_1) = \text{length}(u_3)$, $\text{length}(u_2) = \text{length}(u_4)$ and $(u_1, u_2) \neq (u_3, u_4)$. Then, it leads us to determine if $\max(v_1 \square_9 v_2) > \max(v_3 \square_9 v_4)$ or $\max(v_1 \square_9 v_2) < \max(v_3 \square_9 v_4)$ where $v_1 < v_3$, $v_2 > v_4$. If we consider $v_1 = a$, $v_2 = bb$, $v_3 = ab$ and $v_4 = b$, then we get $\max(v_1 \square_9 v_2) = abb = \max(v_3 \square_9 v_4)$.

By using computation programs realised with **Maxima**, (*c.f.* Section 6) we get:

Lemma 18. *Let n be a positive integer smaller than or equal to 7. Then $K_{n,2} = \{0\}$.*

Proposition 19. *Let X be the alphabet $\{a, b\}$ and \mathcal{S} be the set defined by $\mathcal{S} = \{C_1 \dots C_{10}\}$ equipped with the relation \equiv such that: for any A and B in \mathcal{S} , $A \equiv B$ if and only if there exists an homogenous isomorphism between $(\mathbb{K}\langle X \rangle, \square_A)$ and $(\mathbb{K}\langle X \rangle, \square_B)$ where \square_A (respectively \square_B) is the shuffle product associated to A (respectively B). Let n be the number of isomorphic classes.*

Then $n \in \{7, 8\}$.

2.3 Weak shuffle products on $\mathbb{K}\langle\{a, b, c\}\rangle$

Let $X = \{a, b, c\}$ be an alphabet of cardinality 3. Let C be the 12-tuple $C = (f_1(a \otimes b), f_1(b \otimes a), f_1(b \otimes c), f_1(c \otimes b), f_1(a \otimes c), f_1(c \otimes a), f_1(a \otimes a), f_2(a \otimes a), f_1(b \otimes b), f_2(b \otimes b), f_1(c \otimes c), f_2(c \otimes c))$.

By using Theorem 10, if $(k, m) \in (\mathbb{K}^*)^2$ and $\alpha \in \mathbb{K}$ then C is one of the following tuples

$$\begin{array}{ll}
C_1 = (0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0), & C_2 = (0, 0, 0, 0, 0, 0, 1, 1, 0, 0, 0, 0), \\
C_3 = (0, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0), & C_4 = (0, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1), \\
C_5 = (k, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0), & C_6 = (k, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 1), \\
C_7 = (1, 0, 0, 0, 0, 0, 1, 1, 0, 0, 0, 0), & C_8 = (1, 0, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0), \\
C_9 = (1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0), & C_{10} = (1, 0, 0, 0, 0, 0, 1, 1, 0, 0, 1, 1), \\
C_{11} = (1, 0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 1), & C_{12} = (1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1), \\
C_{13} = (1, 1, 0, 0, 0, 0, 1, 1, 1, 1, 0, 0), & C_{14} = (1, 1, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1), \\
C_{15} = (k, 0, 0, m, 0, 0, 0, 0, 0, 0, 0, 0), & C_{16} = (k, 0, 0, 1, 0, 0, 0, 0, 0, 0, 1, 1), \\
C_{17} = (1, 0, 0, 1, 0, 0, 0, 0, 1, 1, 0, 0), & C_{18} = (1, 0, 0, 1, 0, 0, 1, 1, 1, 1, 0, 0), \\
C_{19} = (1, 0, 0, 1, 0, 0, 1, 1, 0, 0, 1, 1), & C_{20} = (1, 0, 0, 1, 0, 0, 1, 1, 1, 1, 1, 1), \\
C_{21} = (k, 0, 0, 0, m, 0, 0, 0, 0, 0, 0, 0), & C_{22} = (1, 0, 0, 0, 1, 0, 1, 1, 0, 0, 0, 0), \\
C_{23} = (1, 0, 0, 0, 1, 0, \alpha, 1 - \alpha, 0, 0, 0, 0), & C_{24} = (k, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 1), \\
C_{25} = (1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 0, 0), & C_{26} = (1, 0, 0, 0, 1, 0, \alpha, 1 - \alpha, 1, 1, 0, 0), \\
C_{27} = (1, 0, 0, 0, 1, 0, 0, 0, 1, 1, 1, 1), & C_{28} = (1, 0, 0, 0, 1, 0, 1, 1, 1, 1, 1, 1), \\
C_{29} = (1, 0, 0, 0, 1, 0, \alpha, 1 - \alpha, 1, 1, 1, 1), & C_{30} = (k, 0, m, 0, 1, 0, 0, 0, 0, 0, 0, 0), \\
C_{31} = (1, 0, k, 0, 1, 0, 1, 1, 0, 0, 0, 0), & C_{32} = (1, 0, 1, 0, 1, 0, 0, 0, 1, 1, 0, 0), \\
C_{33} = (k, 0, 1, 0, 1, 0, 0, 0, 0, 0, 1, 1), & C_{34} = (1, 0, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0), \\
C_{35} = (1, 0, 1, 0, 1, 0, 1, 1, 0, 0, 1, 1), & C_{36} = (1, 0, 1, 0, 1, 0, 0, 0, 1, 1, 1, 1), \\
C_{37} = (1, 0, 1, 0, 1, 0, 1, 1, 1, 1, 1, 1), & C_{38} = (1, 0, k, 0, 1, 0, \alpha, 1 - \alpha, 0, 0, 0, 0), \\
C_{39} = (1, 0, 1, 0, 1, 0, \alpha, 1 - \alpha, 1, 1, 0, 0), & C_{40} = (1, 0, 1, 0, 1, 0, \alpha, 1 - \alpha, 0, 0, 1, 1), \\
C_{41} = (1, 0, 1, 0, 1, 0, \alpha, 1 - \alpha, 1, 1, 1, 1), & C_{42} = (1, 1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0), \\
C_{43} = (1, 1, 1, 0, 1, 0, 1, 1, 1, 1, 1, 1), & C_{44} = (1, 1, 0, 1, 0, 1, 1, 1, 1, 1, 0, 0),
\end{array}$$

$$C_{45} = (1, 1, 0, 1, 0, 1, 1, 1, 1, 1, 1, 1),$$

$$C_{46} = (1, 1, 0, 1, 0, 1, 1, 1, 1, 1, \alpha, 1 - \alpha),$$

$$C_{47} = (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1).$$

Proposition 20. *Let X be the alphabet $\{a, b, c\}$ and \mathcal{S} be the set defined by $\mathcal{S} = \{C_1 \dots C_{47}\}$ equipped with the relation \equiv such that: for any A and B in \mathcal{S} , $A \equiv B$ if and only if there exists an homogenous isomorphism between $(\mathbb{K}\langle X \rangle, \square_A)$ and $(\mathbb{K}\langle X \rangle, \square_B)$ where \square_A (respectively \square_B) is the shuffle product associated to A (respectively B). Let n be the number of isomorphic classes. Then $n \in \llbracket 33, 39 \rrbracket$.*

Proof. Thanks to Proposition 13, in any set, it is sufficient to consider that $k = m = 1$. Thanks to Proposition 12, we can prove that cases C_{22} and C_{23} are isomorphic, cases C_{25} and C_{26} are isomorphic, cases C_{28} and C_{29} are isomorphic, cases C_{31} and C_{38} are isomorphic, cases C_{34} and C_{39} are isomorphic, cases C_{35} and C_{40} are isomorphic, cases C_{37} and C_{41} are isomorphic and cases C_{45} and C_{46} are isomorphic.

Let K_1 , K_2 and K_3 be the sets defined by:

- $K_1 = \left\{ u = \sum_{x \in X} \lambda_x x, u^2 = 0 \right\},$
- $K_2 = \left\{ u = \sum_{\substack{w \in X^* \\ \text{length}(w)=2}} \lambda_w w, u^2 = 0 \right\},$
- $K_3 = \left\{ u = \sum_{\substack{w \in X^* \\ \text{length}(w)=3}} \lambda_w w, u^2 = 0 \right\}.$

By using K_1 and K_2 , we conclude that C_6 , C_7 and C_8 are in three different isomorphic classes, C_9 , C_{10} and C_{11} are in three different isomorphic classes, C_{16} , C_{17} , C_{22} and C_{24} are in four different isomorphic classes, C_{18} , C_{19} , C_{25} and C_{27} are in four different isomorphic classes, C_{15} and C_{21} are in two different isomorphic classes, C_{31} , C_{32} and C_{33} are in three different isomorphic classes, C_{34} , C_{35} and C_{36} are in three different isomorphic classes, C_{42} and C_{44} are in two different isomorphic classes. With K_3 , we prove that C_{20} and C_{28} are in two different isomorphic classes. Those sets do not enable us to conclude if there exists an isomorphism between C_9 and C_{13} , between C_{12} and C_{14} , between C_{34} and C_{42} , between C_{36} and C_{44} , between C_{43} and C_{47} , between C_{45} and C_{47} . \square

3 Weak shuffle algebras, dendriform algebras, quadri-algebras

Dendriform algebras [22] and quadri-algebras [1] are algebraic structures which enables one to split the associativity. Actually, a dendriform algebra is an algebra \mathcal{A} equipped with a left product \prec and a right product \succ making the couple $(\mathcal{A}, \prec + \succ)$ into an associative algebra and satisfying compatibilities. A quadri-algebra is obtained by splitting each product of a dendriform algebra in two products and the four new products must respect compatibilities. So, a quadri-algebra leads to two dendriform structures and the sum of the four products gives an associative product.

Those two notions have been extensively studied. For instance, Loday and Ronco give the free dendriform algebra on one generator as an algebra over binary planar trees [23]. Thanks to dendriform algebras, Foissy proves [9, proposition 31] that the decorated Hopf algebra of Loday and Ronco and the decorated Hopf algebra of planar rooted trees are isomorphic. Analogue theorems of the Cartier-Quillen-Milnor-Moore theorem have been proved: by Ronco [27] for dendriform algebras, by Chapoton [3] for dendriform bialgebras and by Foissy [10] for bidendriform bialgebras. The bidendriform case implies that **FQSym** is isomorphic to one decorated Hopf algebra of planar rooted trees.

About quadri-algebras, Aguiar and Loday [1] have determined a quadri-algebra structure on infinitesimal algebras and have focused on the free quadri-algebra on one generator. Vallette [32] has proved some conjectures given by Aguiar and Loday in [1, conjectures 4.2, 4.5 and 4.6]. Foissy has presented the free quadri-algebra on one generator as a sub-object of \mathbf{FQSym} [11].

In this section, we recall the dendriform algebra structure and the quadri-algebra structure underlying the classical shuffle algebra. Then, we consider the case of weak shuffle algebras. We prove that they can be equipped with a dendriform structure yet only two weak shuffle products can be considered as coming from a quadri-algebra.

3.1 Dendriform algebras

3.1.1 Background

Definition 21. A dendriform algebra is a vector space \mathcal{D} equipped with two \prec products \succ such that $\forall x, y, z \in \mathcal{D}$,

$$\begin{aligned}(x \prec y) \prec z &= x \prec (y \prec z) + x \prec (y \succ z), \\ (x \succ y) \prec z &= x \succ (y \prec z), \\ (x \prec y) \succ z + (x \succ y) \succ z &= x \succ (y \succ z).\end{aligned}$$

Theorem 22. Let X be a countable alphabet and \sqcup be the classical shuffle product. We define \prec and \succ respectively by:

$$au \prec bv = a(u \sqcup bv), \quad au \succ bv = b(au \sqcup v),$$

for any letters a and b and any words u and v . Then $(\mathbb{K}\langle X \rangle, \prec, \succ)$ is a dendriform algebra and for any words u and v

$$u \sqcup v = u \prec v + u \succ v.$$

Theorem 23. Let X be a countable alphabet and \sqcup be the classical shuffle product. We define \wedge and \vee respectively by:

$$ua \wedge vb = (u \sqcup vb)a, \quad ua \vee vb = (ua \sqcup v)b,$$

for any letters a and b and any words u and v . Then $(\mathbb{K}\langle X \rangle, \wedge, \vee)$ is a dendriform algebra and for any words u and v

$$u \sqcup v = u \wedge v + u \vee v.$$

3.1.2 Weak shuffle products

Theorem 24. Let X be a countable alphabet and \square be a weak shuffle product such that $f_1(a \otimes a) \in \{0, 1\}$ for any letter $a \in X$. We define the products \prec and \succ respectively by:

$$au \prec bv = f_1(a \otimes b)a(u \square bv), \quad au \succ bv = f_2(a \otimes b)b(au \square v),$$

for any letters a and b and any words u and v . Then $(\mathbb{K}\langle X \rangle, \prec, \succ)$ is a dendriform algebra.

Proof. Let \square be a weak shuffle product and let a, b and c be three letters of X . Then:

$$\begin{aligned}(a \prec b) \prec c &= f_1(a \otimes b)f_1(a \otimes c)f_1(b \otimes c)abc + f_1(a \otimes b)f_1(a \otimes c)f_2(b \otimes c)acb, \\ a \prec (b \square c) &= f_1(a \otimes b)f_1(b \otimes c)abc + f_1(a \otimes c)f_2(b \otimes c)acb, \\ (a \succ b) \prec c &= f_2(a \otimes b)f_1(b \otimes c)f_1(a \otimes c)bac + f_2(a \otimes b)f_1(b \otimes c)f_2(a \otimes c)bca, \\ a \succ (b \prec c) &= f_2(a \otimes b)f_1(b \otimes c)f_1(a \otimes c)bac + f_2(a \otimes b)f_1(b \otimes c)f_2(a \otimes c)bca, \\ (a \square b) \succ c &= f_1(a \otimes b)f_2(a \otimes c)cab + f_2(a \otimes b)f_2(b \otimes c)cba, \\ a \succ (b \succ c) &= f_2(b \otimes c)f_2(a \otimes c)f_1(a \otimes b)cab + f_2(b \otimes c)f_2(a \otimes c)f_2(a \otimes b)cba.\end{aligned}$$

Then $(a \succ b) \prec c = a \succ (b \prec c)$. If the three letters are all distinct or only two of them are equal or $a = b = c$ with $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$ the relations given by Theorem 10 imply $(a \prec b) \prec c = a \prec (b \square c)$ and $(a \square b) \succ c = a \succ (b \succ c)$. If $a = b = c$ with $f_1(a \otimes a) = 1 - f_2(a \otimes a)$ then $(a \prec a) \prec a = a \prec (a \square a)$ and $(a \square a) \succ a = a \succ (a \succ a)$ if and only if $f_1(a \otimes a) \in \{0, 1\}$ and then $f_1(a \otimes a)f_2(a \otimes a) = 0$.

We assume now there exists an integer $n \leq 3$ such that, for any non-empty words u, v and w with $\text{length}(u) + \text{length}(v) + \text{length}(w) = n$, relations $(u \prec v) \prec w = u \prec (v \square w)$, $(u \succ v) \prec w = u \succ (v \prec w)$ and $(u \square v) \succ w = u \succ (v \succ w)$ are satisfied.

Let u, v and w be three non-empty words such that $\text{length}(u) + \text{length}(v) + \text{length}(w) = n + 1$. There exist three letters a, b and c , not necessarily distinct and three words u_1, v_1 and w_1 , not necessarily non-empty, such that $u = au_1, v = bv_1$ and $w = cw_1$. Then

1.

$$\begin{aligned} (u \prec v) \prec w &= f_1(a \otimes b)f_1(a \otimes c)a[(u_1 \square bv_1) \square cw_1] = f_1(a \otimes b)f_1(a \otimes c)a[u_1 \square (bv_1 \square cw_1)] \\ &= f_1(a \otimes b)f_1(a \otimes c)f_1(b \otimes c)a[u_1 \square b(v_1 \square cw_1)] \\ &\quad + f_1(a \otimes b)f_1(a \otimes c)f_2(b \otimes c)a[u_1 \square c(bv_1 \square w_1)], \\ u \prec (v \square w) &= f_1(b \otimes c)f_1(a \otimes b)a[u_1 \square b(v_1 \square cw_1)] \\ &\quad + f_2(b \otimes c)f_1(a \otimes c)f_1(a \otimes c)a[u_1 \square c(bv_1 \square w_1)]. \end{aligned}$$

2.

$$\begin{aligned} (u \succ v) \prec w &= f_2(a \otimes b)f_1(b \otimes c)b[(au_1 \square v_1) \square cw_1], \\ u \succ (v \prec w) &= f_1(b \otimes c)f_2(a \otimes b)b[au_1 \square (v_1 \square cw_1)]. \end{aligned}$$

3.

$$\begin{aligned} (u \square v) \succ w &= f_1(a \otimes b)f_2(a \otimes c)c[a(u_1 \square bv_1) \square cw_1] + f_2(a \otimes b)f_2(b \otimes c)c[b(au_1 \square v_1) \square cw_1], \\ u \succ (v \succ w) &= f_2(b \otimes c)f_2(a \otimes c)c[au_1 \square (bv_1 \square w_1)] = f_2(b \otimes c)f_2(a \otimes c)c[(au_1 \square bv_1) \square w_1] \\ &= f_2(b \otimes c)f_2(a \otimes c)f_1(a \otimes b)c[a(u_1 \square bv_1) \square w_1] \\ &\quad + f_2(b \otimes c)f_2(a \otimes c)f_2(a \otimes b)c[b(au_1 \square v_1) \square w_1]. \end{aligned}$$

Thus, $(u \prec v) \prec w = u \prec (v \square w)$, $(u \succ v) \prec w = u \succ (v \prec w)$ and $(u \square v) \succ w = u \succ (v \succ w)$. \square

By considering the right hand side rather than the left hand side, we get the following definition and theorem.

Definition 25. Let X be a countable alphabet. An end weak shuffle product on $\mathbb{K}\langle X \rangle$ is an associative and commutative product \square_E such that for any $(a, b) \in (X)^2$ and any $(u, v) \in (X^*)^2$ then

$$ua \square_E vb = f_{1,E}(a \otimes b)(u \square_E vb)a + f_{2,E}(a \otimes b)(ua \square_E v)b,$$

where $f_{1,E}$ and $f_{2,E}$ are linear maps from $\mathbb{K}.X \otimes \mathbb{K}.X$ to \mathbb{K} , $u \square_E 0 = 0 \square_E u = 0$ and $u \square_E 1 = 1 \square_E u = u$.

Theorem 26. Let X be a countable alphabet and let \square_E be an end weak shuffle product such that $f_{1,E}(a \otimes a) \in \{0, 1\}$ for any letter $a \in X$. We define the products \wedge and \vee by:

$$ua \wedge vb = f_{1,E}(a \otimes b)(u \square_E vb)a, \quad au \vee bv = f_{2,E}(a \otimes b)(ua \square_E v)b,$$

for any letters a and b and any words u and v . Then $(\mathbb{K}\langle X \rangle, \wedge, \vee)$ is a dendriform algebra.

Remark. Let α be a real number. Let \square be the weak shuffle product satisfying $f_1(a \otimes a) = 1 - f_2(a \otimes a) = \alpha$ for a unique letter a . Even if \square does not depend on the value of α , to express the algebra as a dendriform algebra the assumption $\alpha \in \{0, 1\}$ is necessary.

3.2 Quadri-algebras

3.2.1 Background

Definition 27. A quadri-algebra is \mathcal{Q} is a vector space equipped with four products $\searrow, \nearrow, \swarrow$ and \nwarrow such that: for any $x, y, z \in \mathcal{Q}$,

$$\begin{aligned} (x \swarrow y) \searrow z &= x \searrow (y \cdot z), & (x \nearrow y) \searrow z &= x \nearrow (y \prec z), \\ (x \nwarrow y) \searrow z &= x \nwarrow (y \wedge z), & (x \searrow y) \searrow z &= x \searrow (y \swarrow z), \\ (x \prec y) \nwarrow z &= x \nwarrow (y \vee z), & (x \succ y) \nwarrow z &= x \searrow (y \nwarrow z), \end{aligned}$$

and

$$\begin{aligned} (x \wedge y) \nearrow z &= x \nearrow (y \succ z), \\ (x \vee y) \nearrow z &= x \searrow (y \nearrow z), \\ (x \cdot y) \searrow z &= x \searrow (y \searrow z). \end{aligned}$$

where

$$\begin{aligned} x \prec y &= x \swarrow y + x \nwarrow y, & x \wedge y &= x \nearrow y + x \swarrow y, \\ x \succ y &= x \nearrow y + x \searrow y, & x \vee y &= x \searrow y + x \nwarrow y, \end{aligned}$$

and

$$x \cdot y = x \swarrow y + x \nwarrow y + x \nearrow y + x \searrow y = x \prec y + x \succ y = x \wedge y + x \vee y.$$

Theorem 28. Let X be a countable alphabet and let \sqcup be the classical shuffle product. The products $\searrow, \nearrow, \swarrow$ and \nwarrow are defined as follow:

$$\begin{aligned} auc \swarrow bvd &= a(u \sqcup bvd)c, & auc \nwarrow bvd &= a(uc \sqcup bv)d, \\ auc \nearrow bvd &= b(au \sqcup vd)c, & auc \searrow bvd &= b(auc \sqcup v)d \end{aligned}$$

for any letters a, b, c and d and any words u and v . Then $(\mathbb{K}\langle X \rangle, \searrow, \nearrow, \swarrow, \nwarrow)$ is a quadri-algebra.

Proof. It is proved in [1, Section 1.8]. The main ingredient of the proof is the following statement: for any letters a, b, c and d and any words u and v we have

$$auc \sqcup bvd = a(uc \sqcup bvd) + b(auc \sqcup vd) = (au \sqcup bvd)c + (auc \sqcup bv)d.$$

□

3.2.2 Weak shuffle algebras

Proposition 29. Let X be a countable alphabet of cardinality at least 2. Let \square be a weak shuffle product. There exists an end weak shuffle product \square_E such that $\square = \square_E$ if, and only if, \square is the null product or the classical shuffle product.

Proof. It is sufficient to prove the proposition for an alphabet of cardinality 2 and assume images of functions $f_1, f_2, f_{1,E}$ and $f_{2,E}$ are subsets of $\{0, 1\}$. Let C be the 6-tuple $C =$

$$\left(f_1(a \otimes b), f_1(b \otimes a), f_1(a \otimes a), f_2(a \otimes a), f_1(b \otimes b), f_2(b \otimes b) \right).$$

Case $C = (1, 0, 0, 0, 0, 0)$. If $\square = \square_E$ then

$$\begin{aligned} a\square_E ba &= (f_{1,E}(a \otimes a) + f_{2,E}(a \otimes a)f_{1,E}(a \otimes b)) baa + f_{2,E}(a \otimes a)f_{1,E}(b \otimes a)aba \\ &= a\square ba = aba. \end{aligned}$$

Thus $f_{2,E}(a \otimes a) = 1$ and then $a\square_E a = (f_{1,E}(a \otimes a) + 1)aa \neq 0$ and yet $a\square a = 0$. Contradiction.

Cases $C = (1, 0, 1, 1, 0, 0)$ and $C = (1, 0, 1, 0, 0, 0)$. We recall that these two cases are isomorphic. If $\square = \square_E$ then

$$\begin{aligned} a\square_E ba &= (f_{1,E}(a \otimes a) + f_{2,E}(a \otimes a)f_{1,E}(a \otimes b)) baa + f_{2,E}(a \otimes a)f_{1,E}(b \otimes a)aba \\ &= (f_{1,E}(a \otimes a)f_{1,E}(a \otimes b) + f_{2,E}(a \otimes a)) baa + f_{1,E}(a \otimes a)f_{1,E}(b \otimes a)aba \\ &= ba\square_E a = a\square ba = aba. \end{aligned}$$

Thus $f_{1,E}(a \otimes a) = f_{2,E}(a \otimes a) = f_{1,E}(b \otimes a) = 1$ and $f_{1,E}(a \otimes b) = -1$. Contradiction.

Cases $C = (1, 0, 1, 0, 1, 1)$ and $C = (1, 0, 1, 1, 1, 1)$. The same calculations as in the previous case answer the question.

Case $C = (1, 0, 0, 0, 1, 1)$. If $\square = \square_E$ then

$$\begin{aligned} ba\square_E b &= f_{1,E}(a \otimes b) (f_{1,E}(b \otimes b) + f_{2,E}(b \otimes b)) bba + f_{1,E}(b \otimes a)bab \\ &= ba\square b = bba + bab. \end{aligned}$$

Thus $f_{1,E}(a \otimes b) = f_{1,E}(b \otimes a) = f_{1,E}(a \otimes a) = f_{2,E}(a \otimes a) = f_{1,E}(b \otimes b) = f_{2,E}(b \otimes b) = 1$ with $f_{1,E}(b \otimes b) + f_{2,E}(b \otimes b) = 1$. Contradiction.

Cases $C = (0, 0, 1, 1, 0, 0)$. If $\square = \square_E$ then

$$\begin{aligned} ab\square_E a &= f_{1,E}(b \otimes a) (f_{1,E}(a \otimes a) + f_{2,E}(a \otimes a)) aab + f_{1,E}(a \otimes b)aba \\ &= ab\square a = aab. \end{aligned}$$

Thus $f_{1,E}(a \otimes b) = 0$, $f_{1,E}(b \otimes a) = 1$ and $f_{1,E}(a \otimes a) + f_{2,E}(a \otimes a) = 1$. Contradiction.

Cases $C = (0, 0, 1, 1, 1, 1)$. The same calculations as in the previous case answer the question. □

Corollary 30. *The construction used in Theorem 28 does not lead to a quadri-algebra structure on a weak shuffle product \square except if \square is the null shuffle or the classical shuffle.*

4 Relations on weak stuffle products

Proposition 31. *Let X be a countable alphabet, let a, b and c be three distinct letters in X and \square a weak stuffle product. Then:*

1. *By using the maps f_1 and f_2 coming from \square , we define the product \square' by: $au\square'bv = f_1(a \otimes b)a(u\square'bv) + f_2(a \otimes b)b(au\square'v)$ for any letters a and b and any words u and v . The product \square' is a weak shuffle product.*
2. *The function f_3 is associative and commutative.*
3. *If $f_3(a \otimes a) \neq 0$ then $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$.*
4. *If $f_3(a \otimes b) \neq 0$ then $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$ and $f_1(b \otimes b) = f_2(b \otimes b) \in \{0, 1\}$.*

5. If $f_3(a \otimes a) \in \mathbb{K}^*a$ then $f_1(b \otimes a) \in \{0, 1\}$.

6. If $f_3(a \otimes a) \in \mathbb{K}^*b$ then

(a) If $f_3(a \otimes b) \neq 0$ or $f_3(b \otimes b) \neq 0$ or there exists $x \in X \setminus \{a, b\}$ such that $f_3(b \otimes x) \neq 0$ then $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = f_1(a \otimes b) = f_1(b \otimes a) \in \{0, 1\}$.

(b) If $f_3(a \otimes b) = 0$ and $f_3(b \otimes b) = 0$ then

i. either $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = f_1(a \otimes b) = f_1(b \otimes a) \in \{0, 1\}$,
ii. or $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes a) = 1$, $f_1(b \otimes b) + f_2(b \otimes b) = 1$ and $f_1(a \otimes b) = 0$.

(c) For any $x \in X \setminus \{a, b\}$ then

i. $f_1(a \otimes x) = f_1(b \otimes x)$,
ii. $f_1^2(x \otimes a) = f_1(x \otimes b)$.

7. If $f_3(a \otimes b) \in \mathbb{K}^*a$ then:

(a) $f_1(b \otimes a) = f_1(a \otimes a)f_1(a \otimes b) = f_1(b \otimes a)f_1(b \otimes b)$.

(b) $f_1(a \otimes b) = f_1(b \otimes b)$.

(c) For any $x \in X \setminus \{a, b\}$ such that $f_3(b \otimes x) \notin \mathbb{K}^*x$ then

i. $f_1(a \otimes x) = f_1(b \otimes x)$,
ii. $f_1(x \otimes a)[1 - f_1(x \otimes b)] = 0$.

(d) For any $x \in X \setminus \{a, b\}$ such that $f_3(b \otimes x) \in \mathbb{K}^*x$ then

i. $f_1(b \otimes a) = f_1(x \otimes a)f_1(x \otimes b)$,
ii. $f_1(b \otimes x) = f_1(a \otimes b)f_1(a \otimes x)$,

8. If $f_3(a \otimes b) \in \mathbb{K}^*c$ then:

(a) $f_1(c \otimes c) = f_2(c \otimes c) \in \{0, 1\}$.

(b) $f_1(b \otimes a) = f_1(c \otimes a) = f_1(a \otimes a)$.

(c) $f_1(a \otimes b) = f_1(c \otimes b) = f_1(b \otimes b)$.

(d) $f_1(a \otimes c) = f_1(a \otimes a)f_1(b \otimes b) = f_1(b \otimes c) = f_1(c \otimes c)$.

Proof. 1. Let a and b be two letters and let u and v be two words. By using words of length $\text{length}(u) + \text{length}(v) + 2$ appearing in $au \square bv$, we get the statement. In the sequel, the use of the relations given in Theorem 10 is implied.

2. By using words of length 1 appearing in $x \square y$, $x \square y$, $(x \square y) \square z$ and $x \square (y \square z)$ for any letters x, y, z , we prove that the function f_3 is associative and commutative.

3. We assume $f_3(a \otimes a) \neq 0$. Since $a \square aa = aa \square a$ and $(a \square a) \square aa = a \square (a \square aa)$ then $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$.

4. We assume $f_3(a \otimes b) \neq 0$. Since $a \square ab = ab \square a$, $b \square ba = ba \square b$, $(a \square b) \square a = (a \square a) \square b$ and $(b \square a) \square b = (b \square b) \square a$ then $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$ and $f_1(b \otimes b) = f_2(b \otimes b) \in \{0, 1\}$.

5. This item is proved by using $(a \square a) \square b = (a \square b) \square a$ and $a \square (a \square ba) = (a \square a) \square ba$.

6. We assume $f_3(a \otimes a) \in \mathbb{K}^*b$.

(a) If $f_3(a \otimes b) \neq 0$ or $f_3(b \otimes b) \neq 0$, since $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$, $f_1(b \otimes b) = f_2(b \otimes b) \in \{0, 1\}$, $(a \square b) \square a = (a \square a) \square b$ and $(a \square a) \square aa = a \square (a \square aa)$, then $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes b) = f_2(b \otimes b) = f_1(a \otimes b) = f_1(b \otimes a) \in \{0, 1\}$.

(b) If $f_3(a \otimes b) = 0$ and $f_3(b \otimes b) = 0$, since $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$, $(a \square b) \square a = (a \square a) \square b$ and $(a \square a) \square aa = a \square (a \square aa)$ then we prove the relations.

(c) This item is proved thanks to the relation $(a \square b) \square a = (a \square a) \square b$.

7. We assume $f_3(a \otimes b) \in \mathbb{K}^* a$.

(a) This item is proved by using $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$, $f_1(b \otimes b) = f_2(b \otimes b) \in \{0, 1\}$, $(a \square b) \square a = (a \square a) \square b$ and $(b \square a) \square b = (b \square b) \square a$.

(b) By using $(b \square b) \square a = (b \square a) \square b$ and $(a \square b) \square ba = a \square (b \square ba)$ we prove $f_1(a \otimes b) = f_1(b \otimes b)$.

(c) Those two subitems are proven by using $(a \square b) \square x = (a \square x) \square b = (b \square x) \square a$.

(d) Those two subitems are proven by using $(a \square b) \square x = (a \square x) \square b = (b \square x) \square a$.

8. We assume $f_3(a \otimes b) \in \mathbb{K}^* c$. Then $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$ and $f_1(b \otimes b) = f_2(b \otimes b) \in \{0, 1\}$. By using the relations $(a \square b) \square c = (a \square c) \square b = (b \square c) \square a$, $(a \square b) \square b = (b \square b) \square a$, $(b \square a) \square a = (a \square a) \square b$, $(a \square b) \square aa = a \square (b \square aa) = b \square (a \square aa)$ and $(b \square a) \square bb = b \square (a \square bb) = a \square (b \square bb)$ we prove all subitems. □

Examples.

1. The q -shuffle product associated to the Schlesinger-Zudilin model is the weak stuffle product where $f_1(y \otimes p) = f_1(y \otimes y) = f_1(p \otimes p) = f_2(p \otimes p) = 1$, $f_1(p \otimes y) = f_2(y \otimes y) = 0$, $f_3(p \otimes p) = p$, $f_3(y \otimes p) = f_3(y \otimes y) = 0$.

2. The q -shuffle product associated to the Bradley-Zhao model is the weak stuffle product where $f_1(y \otimes p) = f_1(y \otimes \bar{p}) = f_1(p \otimes \bar{p}) = f_1(\bar{p} \otimes p) = f_1(p \otimes p) = f_2(p \otimes p) = f_1(\bar{p} \otimes \bar{p}) = f_2(\bar{p} \otimes \bar{p}) = f_1(y \otimes y) = 1$, $f_1(p \otimes y) = f_1(\bar{p} \otimes y) = f_2(y \otimes y) = 0$, $f_3(p \otimes p) = p$, $f_3(\bar{p} \otimes \bar{p}) = -\bar{p}$, $f_3(y \otimes p) = f_3(y \otimes y) = f_3(y \otimes \bar{p}) = f_3(p \otimes \bar{p}) = 0$.

Corollary 32. *Let $X = \{x_1, \dots, x_n, \dots\}$ be an infinite countable alphabet. We assume \square is a weak stuffle product such that $f_3(x_i \otimes x_j) \in \mathbb{K}^* x_{i+j}$ for any positive integers i and j . Then, the underlying weak shuffle product is either the null shuffle product or the classical shuffle product i.e. ($f_1 \equiv 0$ and $f_2 \equiv 0$) or ($f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any letters a and b).*

Proof. We use an inductive proof. First of all, since $f_3(x_i \otimes x_i) \neq 0$ for any positive integer i , we have $f_1(x_i \otimes x_i) = f_2(x_i \otimes x_i)$. Besides, $f_3(x_1 \otimes x_1) = x_2 \neq x_1$ and $f_3(x_2 \otimes x_2) \neq 0$, so $f_1(x_1 \otimes x_1) = f_2(x_1 \otimes x_1) = f_1(x_2 \otimes x_2) = f_2(x_2 \otimes x_2) = f_1(x_1 \otimes x_2) = f_1(x_2 \otimes x_1) \in \{0, 1\}$.

We assume there exists $n \in \mathbb{N}^*$ such that $n \geq 2$ and $f_1(x_1 \otimes x_1) = f_1(x_1 \otimes x_m)$ for any $m \in \llbracket 1, n \rrbracket$. Then, $f_3(x_1 \otimes x_n) = x_{n+1}$ and $f_1(x_1 \otimes x_{n+1}) = f_1(x_1 \otimes x_1) f_1(x_1 \otimes x_n) = f_1(x_1 \otimes x_1)$. Thus, $f_1(x_1 \otimes x_1) = f_1(x_1 \otimes x_n)$ for any positive integer n .

We assume now there exists $k \in \mathbb{N}^*$ such that $f_1(x_1 \otimes x_1) = f_1(x_i \otimes x_j)$ for any $i \in \llbracket 1, k \rrbracket$ and any positive integer j . For any $i \in \llbracket 1, k \rrbracket$, we know $f_3(x_i \otimes x_{k+1-i}) = x_{k+1}$ so, $f_1(x_{k+1} \otimes x_i) = f_1(x_{k+1-i} \otimes x_i) = f_1(x_1 \otimes x_1)$. Besides, we know

$$f_1(x_{k+1} \otimes x_{k+1}) = f_2(x_{k+1} \otimes x_{k+1}) = f_1(x_1 \otimes x_{k+1}) = f_1(x_1 \otimes x_1).$$

Since $f_3(x_{k+1} \otimes x_1) = x_{k+2}$, we have $f_1(x_{k+1} \otimes x_{k+2}) = f_1(x_1 \otimes x_{k+2}) = f_1(x_1 \otimes x_1)$. We assume there exists a positive integer j such that $f_1(x_{k+1} \otimes x_{k+1+p}) = f_1(x_1 \otimes x_1)$ for any $p \in \llbracket 1, j \rrbracket$. As $f_3(x_{k+1} \otimes x_{j+1}) = x_{k+j+2}$ then

$$f_1(x_{k+1} \otimes x_{k+j+2}) = f_1(x_{k+1} \otimes x_{k+1}) f_1(x_{k+1} \otimes x_{j+1}) = f_1(x_1 \otimes x_1).$$

Finally, ($f_1 \equiv 0$ and $f_2 \equiv 0$) or ($f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any letters a and b). □

By using the commutativity and the associativity of k_3 we have:

Lemma 33. Let $X = \{a, b\}$ be an alphabet of cardinality 2 and let \square be a weak stuffle product. The map f_3 is one of the following:

1. There exists $(\lambda, \mu) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes a) = \lambda b$, $f_3(a \otimes b) = \mu a$ and $f_3(b \otimes b) = \mu b$.
2. There exists $(\lambda, \mu) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes a) = \lambda a$, $f_3(a \otimes b) = \mu a$ and $f_3(b \otimes b) = \frac{\mu^2}{\lambda} a$.
3. There exists $(\lambda, \mu) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes a) = \lambda a$, $f_3(a \otimes b) = \mu a$ and $f_3(b \otimes b) = \mu b$.
4. There exists $(\lambda, \mu) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes a) = 0$, $f_3(a \otimes b) = \mu a$ and $f_3(b \otimes b) = \lambda b$.
5. There exists $(\lambda, \mu) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes a) = \lambda a$, $f_3(a \otimes b) = 0$ and $f_3(b \otimes b) = \mu b$.
6. There exists $\lambda \in \mathbb{K}^*$ such that $f_3(a \otimes a) = \lambda b$, $f_3(a \otimes b) = 0$ and $f_3(b \otimes b) = 0$.
7. There exists $\lambda \in \mathbb{K}^*$ such that $f_3(a \otimes a) = \lambda a$, $f_3(a \otimes b) = 0$ and $f_3(b \otimes b) = 0$.
8. The map f_3 is the null map.

By using Proposition 31 we have:

Proposition 34. Let $X = \{a, b\}$ be an alphabet of cardinality 2 and let \square be a weak stuffle product. In the previous lemma, if f_3 satisfies

1. Item 1 or item 2, then there are two cases:

- $f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any $(a, b) \in X^2$,
- $f_1 \equiv 0$ and $f_2 \equiv 0$.

2. Item 3 or item 4, then there are four cases:

- $f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any $(a, b) \in X^2$,
- $f_1 \equiv 0$ and $f_2 \equiv 0$,
- $f_1(b \otimes a) = f_1(a \otimes b) = f_1(b \otimes b) = f_2(b \otimes b) = 0$ and $f_1(a \otimes a) = f_2(a \otimes a) = 1$,
- $f_1(a \otimes b) = f_1(b \otimes b) = f_2(b \otimes b) = 1$ and $f_1(b \otimes a) = f_1(a \otimes a) = f_2(a \otimes a) = 0$.

3. Item 5, then we have:

- $f_1(a \otimes b) \in \{0, 1\}$,
- $f_1(b \otimes a) \in \{0, 1\}$,
- $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$,
- $f_1(b \otimes b) = f_2(b \otimes b) \in \{0, 1\}$.

4. Item 6, then there are three cases:

- $f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for any $(a, b) \in X^2$,
- $f_1 \equiv 0$ and $f_2 \equiv 0$,
- $f_1(a \otimes a) = f_2(a \otimes a) = f_1(b \otimes a) = 1$, $f_1(a \otimes b) = 0$ and $f_1(b \otimes b) + f_2(b \otimes b) = 1$

5. Item 7, then we have:

- $f_1(b \otimes a) \in \{0, 1\}$,
- $f_1(a \otimes a) = f_2(a \otimes a) \in \{0, 1\}$.

6. Item 8, then we give the answer in Theorem 10.

Lemma 35. Let $X = \{a, b, c\}$ be an alphabet of cardinality 3 and let \square be a weak stuffle product. The map f_3 is one of the following:

20. There exists $(\lambda, \gamma) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes b) = \lambda a$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \gamma c$, $f_3(b \otimes b) = \lambda b$ and $f_3(c \otimes c) = 0$.
21. There exists $(\lambda, \tau) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes b) = \lambda a$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = 0$, $f_3(b \otimes b) = \lambda b$ and $f_3(c \otimes c) = \tau c$.
22. There exists $(\lambda, \gamma, \mu) \in (\mathbb{K}^*)^3$ such that $\gamma\mu = \lambda^2$, $f_3(a \otimes b) = \lambda a$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \gamma a$, $f_3(b \otimes b) = \mu a$ and $f_3(c \otimes c) = 0$.
23. There exists $(\lambda, \gamma, \tau) \in (\mathbb{K}^*)^3$ such that $f_3(a \otimes b) = \lambda a$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \gamma a$, $f_3(b \otimes b) = \lambda b$ and $f_3(c \otimes c) = 0$.
24. There exists $(\lambda, \gamma) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes b) = \lambda a$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \gamma b$, $f_3(b \otimes b) = \lambda b$ and $f_3(c \otimes c) = 0$.
25. There exists $\lambda \in \mathbb{K}^*$ such that $f_3(a \otimes b) = \lambda a$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = 0$, $f_3(b \otimes b) = \lambda b$ and $f_3(c \otimes c) = 0$.
26. There exists $(\lambda, \gamma, \mu) \in (\mathbb{K}^*)^3$ such that $f_3(a \otimes b) = 0$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \lambda a$, $f_3(b \otimes b) = \gamma b$ and $f_3(c \otimes c) = \mu c$.
27. There exists $(\lambda, \gamma) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes b) = 0$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \lambda c$, $f_3(b \otimes b) = \gamma c$ and $f_3(c \otimes c) = 0$.
28. There exists $(\lambda, \gamma) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes b) = 0$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \lambda c$, $f_3(b \otimes b) = \gamma b$ and $f_3(c \otimes c) = 0$.
29. There exists $(\lambda, \gamma) \in (\mathbb{K}^*)^2$ such that $f_3(a \otimes b) = 0$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \lambda a$, $f_3(b \otimes b) = \gamma b$ and $f_3(c \otimes c) = 0$.
30. There exists $\lambda \in \mathbb{K}^*$ such that $f_3(a \otimes b) = 0$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \lambda b$, $f_3(b \otimes b) = 0$ and $f_3(c \otimes c) = 0$.
31. There exists $\lambda \in \mathbb{K}^*$ such that $f_3(a \otimes b) = 0$, $f_3(a \otimes c) = 0$, $f_3(b \otimes c) = 0$, $f_3(a \otimes a) = \lambda a$, $f_3(b \otimes b) = 0$ and $f_3(c \otimes c) = 0$.
32. The map f_3 is the null map.

Proof. We use the fact that the map f_3 is associative and commutative, and then, we get the lemma by direct quite long calculations. \square

Proposition 36. *Let $X = \{a, b, c\}$ be an alphabet of cardinality 3 and let \square be a weak stuffle product. In the previous lemma, if f_3 satisfies one of the items 1, 2, 5, 6, 8, 14, 15 then either $(f_1 \equiv 0$ and $f_2 \equiv 0)$ or $(f_1(a \otimes b) = 1$ and $f_2(a \otimes b) = 1$ for $(a, b) \in X^2)$.*

5 Weak stuffle product and Hopf algebras

If \square is the classical shuffle product or the classical stuffle product then the algebra $(\mathbb{K}\langle X \rangle, \square)$ can be equipped with a compatible coalgebra structure, thanks to the deconcatenation coproduct, which makes it into a Hopf algebra. Are there other weak stuffle products compatible with the deconcatenation? We begin by recalling the Hopf algebra construction for stuffle algebras given in [16, 18, 17]. We then turn to the case of weak stuffle algebras.

Theorem 37. *Let X be a countable alphabet, let $\mathbb{K}\langle X \rangle$ be the vector space generated by words on the alphabet X . We assume there exists at least one product \diamond on $\mathbb{K}\langle X \rangle$ which is commutative and associative. We define the product \star and the coproduct of deconcatenation Δ by:*

$$au \star bv = a(u \star bv) + b(au \star v) + (a \diamond b)(u \star v)$$

and

$$\Delta(w) = \sum_{\substack{(u,v) \in (\mathbb{K}\langle X \rangle)^2, \\ uv=w}} u \otimes v$$

for any letters a and b and any words u, v and w .

Then $(\mathbb{K}\langle X \rangle, \star, \Delta)$ is a Hopf algebra.

Proof. This theorem is proven in [16, 18, 17] by induction and using the filtration given by the length of words. \square

Theorem 38. Let X be a countable alphabet of cardinality $n \in \mathbb{N} \cup \{+\infty\}$ and let \square be a weak shuffle product on $\mathbb{K}\langle X \rangle$. We denote by Δ the deconcatenation coproduct. If Δ respects \square (i.e. if Δ is an algebra morphism) then the underlying weak shuffle product is the classical shuffle product.

Proof. Let \square be a weak shuffle product. We assume the deconcatenation respects \square . Then, for any distinct letters a and b :

$$\begin{aligned} \Delta(a \square a) &= (f_1(a \otimes a) + f_2(a \otimes a)) \Delta(aa) + \Delta(f_3(a \otimes a)) \\ &= (f_1(a \otimes a) + f_2(a \otimes a)) \Delta(aa) + k(a \otimes a) \Delta(g(a \otimes a)) \\ &= (f_1(a \otimes a) + f_2(a \otimes a)) (aa \otimes 1 + a \otimes a + 1 \otimes aa) \\ &\quad + k(a \otimes a) (g(a \otimes a) \otimes 1 + 1 \otimes g(a \otimes a)) \\ &= \Delta(a) \square \Delta(a) \\ &= (f_1(a \otimes a) + f_2(a \otimes a)) (aa \otimes 1 + 1 \otimes aa) + 2a \otimes a \\ &\quad + k(a \otimes a) (g(a \otimes a) \otimes 1 + 1 \otimes g(a \otimes a)), \\ \Delta(a \square b) &= f_1(a \otimes b) \Delta(ab) + f_1(b \otimes a) \Delta(ba) + k(a \otimes b) \Delta(g(a \otimes b)) \\ &= f_1(a \otimes b) (ab \otimes 1 + a \otimes b + 1 \otimes ab) + f_1(b \otimes a) (ba \otimes 1 + b \otimes a + 1 \otimes ba) \\ &\quad + k(a \otimes b) (g(a \otimes b) \otimes 1 + 1 \otimes g(a \otimes b)) \\ &= \Delta(a) \square \Delta(b) = f_1(a \otimes b) (ab \otimes 1 + 1 \otimes ab) + f_1(b \otimes a) (ba \otimes 1 + 1 \otimes ba) + a \otimes b + b \otimes a \\ &\quad + k(a \otimes b) (g(a \otimes b) \otimes 1 + 1 \otimes g(a \otimes b)). \end{aligned}$$

So, $f_1(a \otimes a) = f_2(a \otimes a) = f_1(a \otimes b) = f_1(b \otimes a) = 1$.

The reversal is a particular case of Theorem 37. \square

6 Computation programs

We give computation programs realised to compute the weak shuffle of two words or to prove Lemma 18. In the sequel we assume the alphabet X is the set of integers $\{1, \dots, c\}$ and a word is a list $[i_1, \dots, i_n]$.

We first present a function which computes the weak shuffle product of two words. This function, called `weak_shuffle_product`, takes as entries a list `Rules` which corresponds to the values taken by f_1 and f_2 and two lists `w1` and `w2` which represent the two words to use for computations. We assume

$$\text{Rules} = \left[f_1(1 \otimes 2), \dots, f_1(1 \otimes c), \dots, f_1(c \otimes 1), \dots, f_1(c \otimes c-1), \right. \\ \left. f_1(1 \otimes 1), f_2(1 \otimes 1), \dots, f_1(c \otimes c), f_2(c \otimes c) \right].$$

As exit, the function return a list. Each element of the result is a list of two elements `A` and `B`: `A` is the number of times the word represented by `B` appears in the weak shuffle product of `w1` and `w2`.

```

weak_shuffle_product (Rules ,w1,w2) :=block ( [n1 ,n2 ,u1 ,u2 ,temp ,res , i , j ,
                                             v1a ,v1b ,v2a ,v2b ,P1 ,P2 ,g ,d ,L ,r ,s ,c] ,
/*————— Initialisation of the values of the left side and
                                                    the right side —————*/
g:0 ,
d:0 ,

/*————— Computation of the cardinality of the alphabet. —————*/
r:length (Rules) ,
s:sort (solve (c*(c+1)=r)) ,
c:subst (s [2] ,c) ,

/*————— Message if the variable Rules does not correspond
                                                    to an alphabet. —————*/
if (notequal (c ,floor (c)) or c<1) then print ("erreur") ,

/*————— Computation of the length of words w1 and w2. —————*/
n1:length (w1) ,
n2:length (w2) ,

/*————— We use the commutativity of the weak shuffle product
                                                    to avoid some sub-cases. The word with the smallest length
                                                    is on the left. —————*/
if n1<=n2 then (
    u1:[ [1] ,w1] ,
    u2:[ [1] ,w2]
)
else ( u1:[ [1] ,w2] ,
    u2:[ [1] ,w1] ,
    temp:n1 ,
    n1:n2 ,
    n2:temp
) ,

res:[ [0] ,[]] ,

/*————— We will use a recursive call. —————*/
if equal (n1 ,0) then (
    /*————— Limit case: w1 is the empty word and
                                                    w2 is any word. —————*/
    res:[ [ [1] ,u2 [2]] ]
)
else (
    /*————— We compute the weak shuffle product thanks to the rela-
    tion:  $au(wsp)bv=f1(a\ ot\ b)a(u(wsp)vb)+ f2(a\ ot\ b)b(ua(wsp)v)$ 
    here u and v are words and a and b are letters. —————*/
    v1a:create_list (u1 [2] [i] ,i ,2 ,n1) ,
    v1b:u1 [2] [1] ,
    v2a:create_list (u2 [2] [i] ,i ,2 ,n2) ,
    v2b:u2 [2] [1] ,
    P1:[ ] ,
    P2:[ ] ,

```

```

    /*—— We detemine  $f_1(v1b \setminus ot v2b)$  and  $f_2(v1b \setminus ot v2b)$ . ——*/
if equal(v1b,v2b) then (
    g: Rules [r+2*(-c+v1b)-1],
    d: Rules [r+2*(-c+v1b)]
    ),
if (v1b<v2b) then (
    g: Rules [(v1b-1)*(c-1)+v2b-1],
    d: Rules [(v2b-1)*(c-1)+v1b]
    ),
if (v1b>v2b) then (
    g: Rules [(v1b-1)*(c-1)+v2b],
    d: Rules [(v2b-1)*(c-1)+v1b-1]
    ),

    /*———— Recursive call. —————*/
if g>0 then (
    P1: weak_shuffle_product(Rules,v1a,u2[2]),
    P1: create_list([g*P1[i][1],append([v1b],P1[i][2])],
    i,1,length(P1))
    ),
if d>0 then (
    P2: weak_shuffle_product(Rules,u1[2],v2a),
    P2: create_list([d*P2[i][1],append([v2b],P2[i][2])],
    i,1,length(P2))
    ),
    res:append(P1,P2)
),

    /*—— We rewrite the result for having only one occurence of
    each distinct words. ——*/
L:create_list(res[i][2],i,1,length(res)),
L:unique(L),
res:create_list([ratsimp(sum(if equal(L[i],res[j][2]) then res[j][1]
else 0, j,1, length(res))),L[i]],i,1,length(L)),

    return(res)
);

```

In the sequel, the functions aim at proving if the following statement is true or not for some low n . Let n be a positive integer and let w_1 , w_2 and w be three non-empty words of length n such that $w_1 \leq w_2 \leq w$ and $w_1 < w$. Then $\max(w_1 \square w_2) < \max(w \square w)$? It is trivial for $n = 1$. For $n = 2$, it comes from computations doing in the proof of 16. Thus, those cases are not treated.

The function `words` aims at building all words of length n with an alphabet of cardinality c . It takes as entries the integers n and c and returns a list where each element is a list coresponding to a word. In the result, words are ordered by the ascending order.

```

words(n,c):=block([res,i,j,U],
res:[],
if n=1 then res:create_list([i],i,1,c),
if n>1 then (
    U:words(n-1,c),

```

```

    res:create_list(append(U[i],[j]),j,1,c,i,1,length(U))
  ),
  return(sort(res))
);

```

The function `spectrum_product` aims at determining words appearing in the weak shuffle product of two words `w1` and `w2`. It takes as entries a list `Rules` which gives the rules of computation for the weak shuffle product, an integer `r` which is the length of the list `Rules`, an integer `c` which is the cardinality of the alphabet, and two lists `w1` and `w2` which represent the two words to use for computations.

As exit, the function return a list ordered thanks to the ascending order where each element is a list representing a word appearing in the weak shuffle product of two words `w1` and `w2`.

```

spectrum_product(Rules,r,c,w1,w2):=block([n1,n2,u1,u2,temp,res,i,j,
                                         v1a,v1b,v2a,v2b,P1,P2,g,d],
/*————— Initialisation of the values of
                                     the left side and the right side —————*/
g:0,
d:0,
/*————— Computation of the length of words w1 and w2. —————*/
n1:length(w1),
n2:length(w2),
/*————— We use the commutativity of the weak shuffle product
                                     to avoid some sub-cases. The word with the smallest length
                                     is on the left. —————*/

if n1<=n2 then (
  u1:w1,
  u2:w2
)
else ( u1:w2,
  u2:w1,
  temp:n1,
  n1:n2,
  n2:temp
),
res:[],

/*————— We will use a recursive call. —————*/
if equal(n1,0) then (
  /*—— Limit case: w1 is the empty word and w2 is any word. ——*/
  res:[u2]
)
else (
  /*—— We compute the weak shuffle product thanks to the rela-
  tion:  $au(wsp)bv=f1(a\ ot\ b)a(u(wsp)vb)+f2(a\ ot\ b)b(ua(wsp)v)$ 
  here  $u$  and  $v$  are words and  $a$  and  $b$  are letters. ——*/
  v1a:deleten(u1,1),
  v1b:u1[1],
  v2a:deleten(u2,1),
  v2b:u2[1],
  P1:[],
  P2:[],

```

```

/*———— We detemine  $f_1(v1b \setminus ot v2b)$  and  $f_2(v1b \setminus ot v2b)$ . ———*/
if equal(v1b,v2b) then (
  g: Rules [r+2*(-c+v1b)-1],
  d: Rules [r+2*(-c+v1b)]
),

if (v1b<v2b) then (
  g: Rules [(v1b-1)*(c-1)+v2b-1],
  d: Rules [(v2b-1).(c-1)+v1b]
),
if (v1b>v2b) then (
  g: Rules [(v1b-1)*(c-1)+v2b],
  d: Rules [(v2b-1).(c-1)+v1b-1]
),

/*———— Recursive call. ———*/
if g>0 then (
  P1:spectrum_product(Rules,r,c,v1a,u2),
  P1:create_list(append([v1b],P1[i]),i,1,length(P1))
),
if d>0 then (
  P2:spectrum_product(Rules,r,c,u1,v2a),
  P2:create_list(append([v2b],P2[i]),i,1,length(P2))
),

  res:append(P1,P2)
),

/*———— Words are written once with the ascending order. ———*/
res:sort(unique(res)),
return(res)
);

```

The function `maximum_product` takes as entries a list `Rules` corresponding to the weak shuffle product, an integer r which is the length of `Rules`, an integer c which is the cardinality of the alphabet, an integer n which is the length of words used, a list W which represents the list of words of length n , an integer l which is the length of W , an integer k which is the level of computation. The function returns a list of length $k - 5$. The first one is a list of only one element which is $\max(W[6] \square_9 W[6])$. In the result, the element p with $2 \leq p \leq k - 5$ is a list of two elements A_p and B_p where $A_p = \max(\max(w_1 \square_9 w_2))$ with $w_1 < W[p]$ and $w_2 \leq W[k]$ and $B_p = \max(W[p] \square_9 W[p])$. This function really depends on the weak shuffle product \square_9 .

```

maximum_product(Rules,r,c,n,W,l,k):=block([res,i,P,init],
  res:[],
  if n>1 then (
    /*————  $W[1]=[1, \dots, 1]$ ,  $W[2]=[1, \dots, 1, 2]$ ,
       $W[3]=[1, \dots, 1, 2, 1]$ ,  $W[4]=[1, \dots, 1, 2, 2]$ ,
       $W[5]=[1, \dots, 1, 2, 1, 1]$ ,  $W[6]=[1, \dots, 1, 2, 1, 2]$ ,
      it is enouth to do an initialisation with
       $W[6]$ . ———*/
    if k=6 then (
      init:last(spectrum_product(Rules,r,c,W[6],W[6])),
      res:[[init]]
    )
  )
)

```

```

),
if (k>6 and k<l+1) then (
  /*———— Recursive call. —————*/
  res : maximum_product (Rules , r , c , n , W , l , k-1),

  /*———— Maximum word in res. ————*/
  P : [ last ( sort ( res [ length ( res ) ] ) ) ] ,

  /*—— P is filled in maximum words in W[i](wsp)W[k]
  for i:1 thru k-1 do (
    P : append ( P , [ last ( spectrum_product ( Rules , r , c , W[i] , W[k] ) ) ] ) )
  ),

  /*—— res is filled in a list of two elements:
    the maximum in P and the maximum in W[K](spw)W[k]. ——*/
  res : append ( res , [ [ last ( sort ( P ) ) ,
    last ( spectrum_product ( Rules , r , c , W[k] , W[k] ) ) ] ] )
  )
),
return ( res )
);

```

The function `proof_statement` determines if the statement given at the beginning of the section is proved for words of length n . As entries, it takes a list `Rules` corresponding to the weak shuffle product and an integer corresponding to the length of words used. It returns a boolean. The boolean is true if the statement is satisfied and false if the statement is not satisfied. Since this function uses `maximum_product`, it depends on the weak shuffle product \square .

```

proof_statement ( Rules , n ) := block ( [ res , P , U , i , p , c , r , s , W , l ] ,
  /*———— Computation of the cardinality of the alphabet. ——*/
  r : length ( Rules ) ,
  s : sort ( solve ( c * ( c + 1 ) = r ) ) ,
  c : subst ( s [ 2 ] , c ) ,

  /*———— Message if the variable Rules
    does not correspond to an alphabet. ——*/
  if ( notequal ( c , floor ( c ) ) or c < 1 ) then print ( "erreur " )
  else (
    /*———— Computations. ——*/
    res : true ,
    /*———— Building of words of length n. ——*/
    W : words ( n , c ) ,
    l : length ( W ) ,
    /*———— Building max(w(wsp)w) and max(max(w_1(wsp)w_2)
    with w_K < w and w_K = w. ——*/
    P : maximum_product ( Rules , r , c , n , W , l , l ) ,
    p : length ( P ) ,
    i : 2 ,
    /*———— Checking of the statement at level i. ——*/
    while ( equal ( res , true ) and i < p + 1 ) do (
      if equal ( P [ i ] [ 1 ] , P [ i ] [ 2 ] ) then ( res : false ) ,
      i : i + 1
    )
  )
);

```



```

),
return (res)
);

```

References

- [1] M. AGUIAR AND J.-L. LODAY, *Quadri-algebras*, J. Pure Appl. Algebra, 191 (2004), pp. 205–221.
- [2] D. M. BRADLEY, *Multiple q -zeta values*, Journal of Algebra, 283 (2005), pp. 752 – 798.
- [3] F. CHAPOTON, *Un théorème de Cartier-Milnor-Moore-Quillen pour les bigèbres dendri-formes et les algèbres braces*, J. Pure Appl. Algebra, 168 (2002), pp. 1–18.
- [4] G. DUCHAMP, F. HIVERT, J.-C. NOVELLI, AND J.-Y. THIBON, *Noncommutative symmetric functions VII: Free quasi-symmetric functions revisited*, Ann. Comb., 15 (2011), pp. 655–673.
- [5] G. DUCHAMP, F. HIVERT, AND J.-Y. THIBON, *Noncommutative symmetric functions VI: Free quasi-symmetric functions and related algebras*, Internat. J. Algebra Comput., 12 (2002), pp. 671–717.
- [6] K. EBRAHIMI-FARD AND L. GUO, *Mixable shuffles, quasi-shuffles and Hopf algebras*, Journal of Algebraic Combinatorics, 24 (2006), pp. 83–101.
- [7] K. EBRAHIMI-FARD, D. MANCHON, AND J. SINGER, *Duality and q -multiple zeta values*, Advances in Mathematics, 298 (2016), pp. 254 – 285.
- [8] K. EBRAHIMI-FARD, D. MANCHON, AND J. SINGER, *The Hopf Algebra of q -Multiple Polylogarithms with Non-positive Arguments*, International Mathematics Research Notices, 2017 (2016), pp. 4882–4922.
- [9] L. FOISSY, *Les algèbres de Hopf des arbres enracinés décorés. II*, Bull. Sci. Math., 126 (2002), pp. 249–288.
- [10] ———, *Bidendriform bialgebras, trees and free quasi-symmetric functions*, J. Pure Appl. Algebra, 209 (2007), pp. 439–459.
- [11] ———, *Free quadri-algebras and dual quadri-algebras*, ArXiv e-prints, (2015).
- [12] L. FOISSY, F. PATRAS, AND J.-Y. THIBON, *Deformations of shuffles and quasi-shuffles*, Annales de l’Institut Fourier, 66 (2016), pp. 209–237.
- [13] I. M. GESSEL, *Multipartite p -partitions and inner products of skew Schur functions*, in Combinatorics and algebra (Boulder, Colo., 1983), vol. 34 of Contemp. Math., Amer. Math. Soc., Providence, RI, 1984, pp. 289–317.
- [14] L. GUO AND W. KEIGHER, *Baxter Algebras and Shuffle Products*, Advances in Mathematics, 150 (2000), pp. 117 – 149.
- [15] M. E. HOFFMAN, *The algebra of multiple harmonic series*, J. Algebra, 194 (1997), pp. 477–495.
- [16] ———, *Quasi-shuffle products*, J. Algebraic Combin., 11 (2000), pp. 49–68.
- [17] ———, *Quasi-shuffle algebras and applications*, arXiv e-prints, (2018), p. arXiv:1805.12464.

- [18] M. E. HOFFMAN AND K. IHARA, *Quasi-shuffle products revisited*, J. Algebra, 481 (2017), pp. 293–326.
- [19] M. E. HOFFMAN AND Y. OHNO, *Relations of multiple zeta values and their algebraic expression*, Journal of Algebra, 262 (2003), pp. 332 – 347.
- [20] R.-Q. JIAN, *Quantum quasi-shuffle algebras II*, Journal of Algebra, 472 (2017), pp. 480 – 506.
- [21] R.-Q. JIAN, M. ROSSO, AND J. ZHANG, *Quantum Quasi-Shuffle Algebras*, Letters in Mathematical Physics, 92 (2010), pp. 1–16.
- [22] J.-L. LODAY, *Dialgebras*, in Dialgebras and related operads, vol. 1763 of Lecture Notes in Math., Springer, Berlin, 2001, pp. 7–66.
- [23] J.-L. LODAY AND M. RONCO, *Hopf algebra of the planar binary trees*, Adv. Math., 139 (1998), pp. 293–309.
- [24] C. MALVENUTO, *Produits et coproduits des fonctions quasi-symétriques et de l’algèbre des descentes*, PhD thesis, Université du Québec à Montréal, Laboratoire de Combinatoire et d’Informatique Mathématique, 1994.
- [25] C. MALVENUTO AND C. REUTENAUER, *Duality between quasi-symmetric functions and the Solomon descent algebra*, J. Algebra, 177 (1995), pp. 967–982.
- [26] C. MAMMEZ, *A propos de l’algèbre de Hopf des mots tassés WMat*, Bulletin des Sciences Mathématiques, (2018).
- [27] M. RONCO, *A Milnor-Moore theorem for dendriform Hopf algebras*, C. R. Acad. Sci. Paris Sér. I Math., 332 (2001), pp. 109–114.
- [28] K.-G. SCHLESINGER, *Some remarks on q -deformed multiple polylogarithms*, arXiv Mathematics e-prints, (2001), p. math/0111022.
- [29] J. SINGER, *On q -analogues of multiple zeta values*, Funct. Approx. Comment. Math., 53 (2015), pp. 135–165.
- [30] J. SINGER, *On Bradley’s q -MZVs and a generalized Euler decomposition formula*, Journal of Algebra, 454 (2016), pp. 92 – 122.
- [31] J. SINGER, *q -Analogues of multiple zeta values and their application in renormalization*, PhD thesis, Der Naturwissenschaftlichen Fakultät, der Friedrich-Alexander-Universität, 2016.
- [32] B. VALLETTE, *Manin products, Koszul duality, Loday algebras and Deligne conjecture*, J. Reine Angew. Math., 620 (2008), pp. 105–164.
- [33] Y. VARGAS, *Hopf algebra of permutation pattern functions*, in 26th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2014), L. J. Billera and I. Novik, eds., vol. AT of DMTCS Proceedings, Chicago, United States, 2014, Discrete Mathematics and Theoretical Computer Science, pp. 839–850.
- [34] D. ZAGIER, *Values of zeta functions and their applications*, in First European Congress of Mathematics, Vol. II (Paris, 1992), vol. 120 of Progr. Math., Birkhäuser, Basel, 1994, pp. 497–512.
- [35] J. ZHAO, *Multiple q -zeta functions and multiple q -polylogarithms*, The Ramanujan Journal, 14 (2007), pp. 189–221.

- [36] W. ZUDILIN, *Algebraic relations for multiple zeta values*, Russian Mathematical Surveys, 58 (2003), pp. 1–29.